

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|  |   |  |
|--|---|--|
| <b>(51) International Patent Classification <sup>6</sup>:</b><br><b>A61K 31/18, 31/335, 31/47, C07C 311/10, C07D 215/12</b>  | <b>A1</b>   | <b>(11) International Publication Number:</b> <b>WO 97/19682</b><br><b>(43) International Publication Date:</b> 5 June 1997 (05.06.97) |
| <b>(21) International Application Number:</b> PCT/US96/19085<br><b>(22) International Filing Date:</b> 27 November 1996 (27.11.96)<br><b>(30) Priority Data:</b><br>08/566,104 1 December 1995 (01.12.95) US<br><b>(60) Parent Application or Grant</b><br>(63) Related by Continuation<br>US 08/566,104 (CIP)<br>Filed on 1 December 1995 (01.12.95)<br><b>(71) Applicant (for all designated States except US):</b> SYNAPTIC PHARMACEUTICAL CORPORATION [US/US]; 215 College Road, Paramus, NJ 07652 (US).<br><b>(72) Inventors; and</b><br><b>(75) Inventors/Applicants (for US only):</b> ISLAM, Imadul [IN/US]; 318 Boatwain Court, Hercules, CA 94547 (US). DHANOA, Daljit, S. [US/US]; 13794 Boquita Drive, Del Mar, CA 92014 (US). FINN, John, M. [US/US]; 38 Camelot Drive, West Milford, NJ 07480 (US). DU, Ping [CN/US]; 427 Poets Way, Mahwah, NJ 07430 (US). GLUCHOWSKI, Charles [US/US]; 153 Chopin Drive, Wayne, NJ 07470 (US). JEON, | <b>(74) Agent:</b> WHITE, John, P.; Cooper & Dunham L.L.P., 1185 Avenue of the Americas, New York, NY 10036 (US).<br><b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).<br><b>Published</b><br><i>With international search report.</i><br><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |  |
| <b>(54) Title:</b> ARYL SULFONAMIDE AND SULFAMIDE DERIVATIVES AND USES THEREOF<br><b>(57) Abstract</b><br><p>This invention is directed to novel aryl sulfonamide and sulfamide compounds which bind selectively to and inhibit the activity of the human Y5 receptor. This invention is also related to uses of these compounds for the treatment of feeding disorders such as obesity, anorexia nervosa, bulimia nervosa, and abnormal conditions such as sexual/reproductive disorders, depression, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure or sleep disturbances and for the treatment of any disease in which antagonism of a Y5 receptor may be useful.</p>   |   |  |

20329  
#13

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |  |    |                          |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia                  | GB | United Kingdom                           | MW | Malawi                   |
| AT | Austria                  | GE | Georgia                                  | MX | Mexico                   |
| AU | Australia                | GN | Guinea                                   | NE | Niger                    |
| BB | Barbados                 | GR | Greece                                   | NL | Netherlands              |
| BE | Belgium                  | HU | Hungary                                  | NO | Norway                   |
| BF | Burkina Faso             | IE | Ireland                                  | NZ | New Zealand              |
| BG | Bulgaria                 | IT | Italy                                    | PL | Poland                   |
| BJ | Benin                    | JP | Japan                                    | PT | Portugal                 |
| BR | Brazil                   | KE | Kenya                                    | RO | Romania                  |
| BY | Belarus                  | KG | Kyrgyzstan                               | RU | Russian Federation       |
| CA | Canada                   | KP | Democratic People's Republic<br>of Korea | SD | Sudan                    |
| CF | Central African Republic | KR | Republic of Korea                        | SE | Sweden                   |
| CG | Congo                    | KZ | Kazakhstan                               | SG | Singapore                |
| CH | Switzerland              | LJ | Liechtenstein                            | SI | Slovenia                 |
| CI | Côte d'Ivoire            | LK | Sri Lanka                                | SK | Slovakia                 |
| CM | Cameroon                 | LR | Liberia                                  | SN | Senegal                  |
| CN | China                    | LT | Lithuania                                | SZ | Swaziland                |
| CS | Czechoslovakia           | LU | Luxembourg                               | TD | Chad                     |
| CZ | Czech Republic           | LV | Latvia                                   | TG | Togo                     |
| DE | Germany                  | MC | Monaco                                   | TJ | Tajikistan               |
| DK | Denmark                  | MD | Republic of Moldova                      | TT | Trinidad and Tobago      |
| EE | Estonia                  | MG | Madagascar                               | UA | Ukraine                  |
| ES | Spain                    | ML | Mali                                     | UG | Uganda                   |
| FI | Finland                  | MN | Mongolia                                 | US | United States of America |
| FR | France                   | MR | Mauritania                               | UZ | Uzbekistan               |
| GA | Gabon                    |    |  | VN | Viet Nam                 |

Aryl Sulfonamide and Sulfamide Derivatives  
and Uses Thereof

Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be found at the end of this application, preceding the sequence listing and the claims.

Background of the Invention

The peptide neurotransmitter neuropeptide Y (NPY) is a 36 amino acid member of the pancreatic polypeptide family with widespread distribution throughout the mammalian nervous system (Dumont et al., 1992). The family includes the namesake pancreatic polypeptide (PP), synthesized primarily by endocrine cells in the pancreas; peptide YY (PYY), synthesized primarily by endocrine cells in the gut; and NPY, synthesized primarily in neurons (Michel, 1991; Dumont et al., 1992; Wahlestedt and Reis, 1993). All pancreatic polypeptide family members share a compact structure involving a "PP-fold" and a conserved C-terminal hexapeptide ending in Tyr<sup>36</sup> (or Y<sup>36</sup> in the single letter code). The striking conservation of Y<sup>36</sup> has prompted the reference to the pancreatic polypeptides' receptors as "Y-type" receptors (Wahlestedt et al., 1987), all of which are proposed to function as seven transmembrane-spanning G protein-coupled receptors (Dumont et al., 1992).

NPY and its relatives elicit a broad range of physiological effects through activation of at least five G protein-coupled receptor subtypes known as Y1, Y2, Y3, Y4 (or PP), and the "atypical Y1". While the Y1, Y2, Y3, and Y4 (or PP) receptors were each described previously in

-2-

both radioligand binding and functional assays, the "atypical Y1" receptor is unique in that its classification is based solely on feeding behavior induced by various peptides including NPY.

5  
The role of NPY in normal and abnormal eating behavior, and the ability to interfere with NPY-dependent pathways as a means to appetite and weight control, are areas of great interest in pharmacological and pharmaceutical  
10 research (Sahu and Kalra, 1993; Dryden et al., 1994). NPY is considered to be the most powerful stimulant of feeding behavior yet described (Clark et al., 1984; Levine and Morley, 1984; Stanley and Leibowitz, 1984). The stimulation of feeding behavior by NPY is thought to occur  
15 primarily through activation of the hypothalamic "atypical Y1" receptor. For example, direct injection of NPY into the hypothalamus of satiated rats can increase food intake up to 10-fold over a 4-hour period (Stanley et al., 1992). Similar studies using other peptides has resulted in a  
20 pharmacologic profile for the "atypical Y1" receptor according to the rank order of potencies of peptides in stimulating feeding behavior as follows:  $NPY_{2-36} \geq NPY \sim PYY \sim [Leu^{31}, Pro^{34}]NPY > NPY_{13-36}$  (Kalra et al., 1991; Stanley et al., 1992). The profile is similar to that of  
25 a Y1-like receptor except for the anomalous ability of  $NPY_{2-36}$  to stimulate food intake with potency equivalent or better than that of NPY. A subsequent report in J. Med. Chem. by Balasubramaniam and co-workers (1994) showed that feeding can be regulated by  $[D-Trp^{32}]NPY$ . While this  
30 peptide was presented as an NPY antagonist, the published data at least in part support a stimulatory effect of  $[D-Trp^{32}]NPY$  on feeding. In contrast to other NPY receptor subtypes, the "feeding" receptor has never been characterized for peptide binding affinity in radioligand  
35 binding assays. The fact that a single receptor could be

-3-

responsible for the feeding response has been impossible to validate in the absence of an isolated receptor protein; the possibility exists, for example, that the feeding response could be a composite profile of Y1 and Y2 subtypes.

This problem has been addressed by cloning rat and human cDNAs which encode a single receptor protein, referred to herein as Y5, whose pharmacologic profile links it to the "atypical Y1" receptor. The identification and characterization by applicants of a single molecular entity which explains the "atypical Y1" receptor allows the design of selective drugs which modulate feeding behavior. It is important to note, though, that any credible means of studying or modifying NPY-dependent feeding behavior must necessarily be highly selective, as NPY interacts with multiple receptor subtypes, as noted above (Dumont et al., 1992).

As used in this invention, the term "antagonist" refers to a compound which decreases the activity of a receptor. In the case of a G-protein coupled receptor, activation may be measured using any appropriate second messenger system which is coupled to the receptor in a cell or tissue in which the receptor is expressed. Some specific but by no means limiting examples of well-known second messenger systems are adenylate cyclase, intracellular calcium mobilization, ion channel activation, guanylate cyclase, and inositol phospholipid hydrolysis. Conversely, the term "agonist" refers to a compound which increases the activity of a receptor.

In order to test compounds for selective binding to the human Y5 receptor the cloned cDNAs encoding both the human and rat Y2 and Y4 (or PP) receptors have been used. The

-4-

human and rat Y5 receptors were disclosed in PCT International Application No. PCT/US95/15646, published June 6, 1996, and filed as a continuation in part of U.S. Serial No. 08/349,025, filed December 2, 1994, the contents of which are hereby incorporated by reference into this application. The human and rat Y2 receptors were disclosed in PCT International Application US95/01469, published August 10, 1995, as WO 95/21245, and filed as a continuation-in-part of U.S. 08/192,288, filed February 3, 1994, the contents of which are hereby incorporated by reference into this application. The human and rat Y4 receptors were disclosed in PCT International Application PCT/US94/14436, published July 6, 1995, as WO 95/17906, and filed as a continuation-in-part of U.S. 08/176,412, filed December 28, 1993, the contents of which are hereby incorporated by reference into this application. The Y1 receptor has been cloned from a variety of species including human, rat and mouse (Larhammar et al, 1992; Herzog et al, 1992; Eva et al, 1990; Eva et al, 1992).

The synthesis of novel aryl sulfonamide and sulfamide compounds are disclosed which bind selectively to the cloned human Y5 receptor compared to the other cloned human NPY receptors, and inhibit the activation of the cloned human Y5 receptor as measured in in vitro assays. The in vitro receptor binding and activation assays described hereinafter were performed using various cultured cell lines, each transfected with and expressing only a single Y-type receptor. In addition, the compounds of the present invention were shown to inhibit in animals either NPY-induced feeding behavior or feeding behavior exhibited by food-deprived animals.

-5-

This invention is also directed to the treatment of feeding disorders such as obesity and bulimia nervosa using the compounds described herein. In addition, the compounds of the present invention may also be used to

5 treat abnormal conditions such as sexual/reproductive disorders, depression, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure or sleep disturbances, or any condition in which antagonism of a Y5 receptor may be useful.


Summary of the Invention

This invention is directed to novel aryl sulfonamide and sulfamide compounds which bind selectively to and inhibit the activity of the human Y5 receptor. This invention is also related to uses of these compounds for the treatment of feeding disorders such as obesity, anorexia nervosa, bulimia nervosa, and abnormal conditions such as sexual/reproductive disorders, depression, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure or sleep disturbances and for the treatment of any disease in which antagonism of a Y5 receptor may be useful.




Detailed Description of the Invention

The present invention is directed to compounds having the structures:



wherein Ar is



wherein each Z is independently N or C;

15

wherein each Y is independently N or C;

wherein p is an integer from 0 to 2;

-8-

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

wherein V is S, O, N, or NR<sub>5</sub>;

5

wherein X is a single bond or -NH-;

wherein each R<sub>2</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>2</sub>-C<sub>4</sub> alkenyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>; phenoxy; phenyl; pyridyl; thiophenyl; naphthyl; phthalimide; C<sub>5</sub>-C<sub>7</sub> lactam, C<sub>5</sub>-C<sub>7</sub> cyclic imide, C<sub>5</sub>-C<sub>7</sub> cyclic amino; wherein the phthalimide, lactam, cyclic imide, or cyclic amine is linked by nitrogen; and wherein the phenoxy, phenyl, pyridyl, thiophenyl, naphthyl, phthalimide, lactam, cyclic imide, or cyclic amine is substituted with H, F, Cl, Br, I, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, or NO<sub>2</sub>;

20

wherein each R<sub>3</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>2</sub>-C<sub>4</sub> alkenyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>; or R<sub>3</sub> and R<sub>4</sub> present on adjacent carbon atoms can constitute C<sub>5</sub>-C<sub>7</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> heterocycloalkyl or C<sub>5</sub>-C<sub>7</sub> heteroaryl;

25

wherein each R<sub>4</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;


30

- 9 -

wherein each  $R_5$  is independently H;  $C_1-C_3$  alkyl;  $C_1-C_3$  monohaloalkyl; or  $C_1-C_3$  polyhaloalkyl;

wherein  $L'$  is  $-NR_1-L-$  or


5




10

wherein  $L$  is  $C_3-C_9$  alkyl;  $C_3-C_9$  alkenyl;  $C_3-C_9$  alkynyl;


15



20




25



30

or



wherein  $R_1$  is H; or  $C_1$ - $C_3$  straight chained alkyl;

5

wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

10

wherein one dashed line is a double bond and the other dashed line is a single bond;

15

wherein each  $R_i$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_5$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_5$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

20

wherein  $i$  is an integer from 1 to 4; wherein  $n$  is an integer from 0 to 3; wherein  $m$  is an integer from 0 to 3;

wherein  $K$  is  $-CH_2-NR_{10}-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-(CH_2)_3-$ ;  $-CH_2-NH-CO-NH-(CH_2)_3-$ ;  $-CO-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-(CH_2)_3-$ ;  $-CH_2-NH-CS-NH-(CH_2)_3-$ ;  $-CS-NH-$

-11-


$\text{CHR}_7 - (\text{CH}_2)_j - ; \quad -\text{CH}_2 - \text{NR}_{10} - \text{CS} - \text{CHR}_7 - (\text{CH}_2)_j ; \quad \text{or} \quad -\text{CH}_2 - \text{N} = \text{CSR}_1 - \text{NH} - (\text{CH}_2)_j ;$

wherein  $j$  is an integer from 0 to 3;


5

wherein  $R_7$  is H;  $\text{C}_1 - \text{C}_6$  alkyl;  $\text{CH}_2\text{OR}_5$ ;  $-(\text{CH}_2)_f\text{NHCO}_2\text{R}_6$ ;  $(\text{CH}_2)_p\text{NHSO}_2\text{R}_5$ ;  $\text{CH}_2\text{N}(\text{R}_{11})_2$ ; phenyl; pyridyl; thiophenyl; or naphthyl;

wherein W is



-13-



wherein Q is O; S; N; NR<sub>9</sub>; or C(R<sub>5</sub>)<sub>2</sub>;

5 wherein b is an integer from 1 to 2;

wherein R<sub>8</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; =O; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl;  
 10 N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>4</sub>; NHSO<sub>2</sub>R<sub>4</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;

wherein R<sub>9</sub> is H; C<sub>1</sub>-C<sub>3</sub> alkyl; COR<sub>5</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>;

15 wherein R<sub>10</sub> is H; or C<sub>1</sub>-C<sub>6</sub> alkyl;


wherein R<sub>11</sub> is H; COR<sub>5</sub>; COR<sub>12</sub>; SO<sub>2</sub>R<sub>5</sub>; SO<sub>2</sub>R<sub>12</sub>; and

wherein  $R_{12}$  is phenoxy; phenyl, pyridyl; thiophenyl; or naphthyl; wherein the phenoxy, phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$  alkylthio,  $NO_2$ , phenyl, pyridyl or thiophenyl; or a pharmaceutically acceptable salt thereof.

The invention also provides for the (+) and (-) enantiomers of the compounds described herein.

In one embodiment the invention provides for a compound as described above, where  $R_1$  is H;

where L is selected from  $C_3-C_9$  alkyl or




where the alkyl is substituted with H,  $OR_5$ , CN,  $C_1-C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl; and




-15-

where W is




In other embodiments of the present invention, the compounds may have the structures where Ar is selected from:



where each of  $R_2$ ,  $R_3$ , and  $R_4$  is independently H; F, Cl, Br or I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl; or  $\text{N}(\text{R}_5)_2$ ; where X is  
20 a single bond; where each  $\text{R}_5$  is independently  $\text{C}_1\text{-C}_3$  alkyl;


where L is selected from  $\text{C}_5\text{-alkyl}$  or  $\text{C}_7\text{-alkyl}$ ;




where  $\text{R}_1$  is H;  $\text{CH}_2\text{OH}$ ; or  $\text{CH}_2\text{OR}_6$ ;

-17-


where W is

and where  $R_9$  is H; or  $C_1$ - $C_4$  alkyl.

5 In other embodiments of the present invention Ar is selected from:




L is



and K is  $-\text{CH}_2-\text{NR}_{10}-\text{CHR}_7-(\text{CH}_2)_j-$ .


Additional embodiments of the present invention include the compounds selected from the group consisting of:

5




10

15




20

25




30




35


5



10




15



20


25




30

35


5



10



15




20


25

30

35




or



15

Additional embodiments of the present invention include those in which L is C<sub>5</sub>-alkyl or C<sub>7</sub>-alkyl.


20 In an embodiment of the invention the compounds have the structure:



30

or

35




In one embodiment of the invention K is  
 $-\text{CH}_2-\text{NR}_{10}-\text{CO}-(\text{CH}_2)_7-$ .

5

In another embodiment of the invention the compound has  
 the structure:

10




15

In yet another embodiment of the present invention K is  
 $-\text{CH}_2-\text{NH}-\text{CO}-\text{NH}-(\text{CH}_2)_7-$ .




In a further embodiment of the invention the compound has the structure:




10

The invention also provides for a method of modifying feeding behavior of a subject which comprises administering to the subject an amount of a compound effective to decrease the consumption of food by the subject so as to thereby modify feeding behavior of the subject, where the compound has the structure:

15



wherein Ar is



5 wherein each Z is independently N or C;

wherein each Y is independently N or C;

wherein p is an integer from 0 to 2;

10

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

wherein V is S, O, N, or  $NR_6$ ;

15

wherein X is a single bond or  $-NH-$ ;

wherein each  $R_2$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; phenoxy; phenyl; pyridyl; thiophenyl; naphthyl; phthalimide;  $\text{C}_5\text{-C}_7$  lactam,  $\text{C}_5\text{-C}_7$  cyclic imide,  $\text{C}_5\text{-C}_7$  cyclic amino; wherein the phthalimide, lactam, cyclic imide, or cyclic amine is linked by nitrogen; and wherein the phenoxy, phenyl, pyridyl, thiophenyl, naphthyl, phthalimide, lactam, cyclic imide, or cyclic amine is substituted with H, F, Cl, Br, I,  $\text{CF}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{C}_1\text{-C}_4$  alkylthio, or  $\text{NO}_2$ ;


wherein each  $R_3$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; or  $\text{R}_2$  and  $\text{R}_3$  present on adjacent carbon atoms can constitute  $\text{C}_5\text{-C}_7$  cycloalkyl,  $\text{C}_5\text{-C}_7$  heterocycloalkyl or  $\text{C}_5\text{-C}_7$  heteroaryl;

wherein each  $R_4$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ; or  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;

wherein each  $R_4$  is independently H;  $\text{C}_1\text{-C}_3$  alkyl;  $\text{C}_1\text{-C}_3$  monohaloalkyl; or  $\text{C}_1\text{-C}_3$  polyhaloalkyl;

wherein  $\text{L}'$  is  $-\text{NR}_1\text{-L-}$  or


- 26 -




5

wherein L is C<sub>3</sub>-C<sub>9</sub> alkyl; C<sub>3</sub>-C<sub>9</sub> alkenyl; C<sub>3</sub>-C<sub>9</sub> alkynyl;

10




15



20

25




30

or

35

-27-



wherein  $R_1$  is H; or  $C_1$ - $C_3$  straight chained alkyl;

15 wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

wherein one dashed line is a double bond and the other dashed line is a single bond;

20

wherein each  $R_6$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_5$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_5$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

25

wherein  $i$  is an integer from 1 to 4; wherein  $n$  is an integer from 0 to 3; wherein  $m$  is an integer from 0 to 3;

30

wherein  $K$  is  $-CH_2-NR_{10}-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-(CH_2)_3-$ ;  $-CH_2-NH-CO-NH-(CH_2)_3-$ ;  $-CO-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-(CH_2)_3-$ ;  $-CH_2-NH-CS-NH-(CH_2)_3-$ ;  $-CS-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-CHR_7-(CH_2)_3-$ ; or  $-CH_2-N=CSR_1-NH-(CH_2)_3-$ ;


35


-28-

wherein  $j$  is an integer from 0 to 3;

wherein  $R_7$  is H;  $C_1$ - $C_6$  alkyl;  $CH_2OR_5$ ;  $(CH_2)_fNHCO_2R_5$ ;  
5  $(CH_2)_pNHSO_2R_5$ ;  $CH_2N(R_{11})_2$ ; phenyl; pyridyl; thiophenyl; or  
naphthyl;

wherein W is





wherein Q is O; S; N; NR<sub>9</sub>; or C(R<sub>5</sub>)<sub>2</sub>;

5 wherein b is an integer from 1 to 2;

wherein R<sub>8</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; =O; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl;  
 10 N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;

wherein R<sub>9</sub> is H; C<sub>1</sub>-C<sub>6</sub> alkyl; COR<sub>5</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>;

15 wherein R<sub>10</sub> is H; or C<sub>1</sub>-C<sub>6</sub> alkyl;



wherein R<sub>11</sub> is H; COR<sub>5</sub>; COR<sub>12</sub>; SO<sub>2</sub>R<sub>6</sub>; SO<sub>2</sub>R<sub>12</sub>; and




wherein R<sub>12</sub> is phenoxy; phenyl, pyridyl; thiophenyl; or naphthyl; wherein the phenoxy, phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, NO<sub>2</sub>, phenyl, pyridyl or thiophenyl; or a pharmaceutically acceptable salt thereof.

In one embodiment of the method described above the subject is a vertebrate, a mammal, a human or a canine. In another embodiment the compound is administered in combination with food.


The invention also provides for a method of modifying feeding behavior where the compound has the structure:




5



10




15



20


25




30

35

5




10




15

20




25

30




35

5




10

15



20


25




30

The invention further provides a method of treating a feeding disorder in a subject which comprises administering to the subject an amount of a compound effective to decrease consumption of food by the subject, where the compound has the structure:

35



wherein Ar is



5 wherein each Z is independently N or C;

wherein each Y is independently N or C;

wherein p is an integer from 0 to 2;

10

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

wherein V is S, O, N, or NR<sub>5</sub>;

15

wherein X is a single bond or -NH-;

-36-

wherein each  $R_2$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1$ - $\text{C}_4$  alkyl;  $\text{C}_2$ - $\text{C}_4$  alkenyl;  $\text{C}_1$ - $\text{C}_4$  alkoxy;  $\text{C}_1$ - $\text{C}_4$  hydroxyalkyl;  $\text{C}_1$ - $\text{C}_4$  methoxyalkyl;  $\text{C}_1$ - $\text{C}_4$  monohaloalkyl;  $\text{C}_1$ - $\text{C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; phenoxy; phenyl; pyridyl; thiophenyl; naphthyl; phthalimide;  $\text{C}_5$ - $\text{C}_7$  lactam,  $\text{C}_5$ - $\text{C}_7$  cyclic imide,  $\text{C}_5$ - $\text{C}_7$  cyclic amino; wherein the phthalimide, lactam, cyclic imide, or cyclic amine is linked by nitrogen; and wherein the phenoxy, phenyl, pyridyl, thiophenyl, naphthyl, phthalimide, lactam, cyclic imide, or cyclic amine is substituted with H, F, Cl, Br, I,  $\text{CF}_3$ ,  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkoxy,  $\text{C}_1$ - $\text{C}_4$  alkylthio, or  $\text{NO}_2$ ;


wherein each  $R_3$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1$ - $\text{C}_4$  alkyl;  $\text{C}_2$ - $\text{C}_4$  alkenyl;  $\text{C}_1$ - $\text{C}_4$  alkoxy;  $\text{C}_1$ - $\text{C}_4$  hydroxyalkyl;  $\text{C}_1$ - $\text{C}_4$  methoxyalkyl;  $\text{C}_1$ - $\text{C}_4$  monohaloalkyl;  $\text{C}_1$ - $\text{C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; or  $\text{R}_2$  and  $\text{R}_3$  present on adjacent carbon atoms can constitute  $\text{C}_5$ - $\text{C}_7$  cycloalkyl,  $\text{C}_5$ - $\text{C}_7$  heterocycloalkyl or  $\text{C}_5$ - $\text{C}_7$  heteroaryl;

wherein each  $R_4$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1$ - $\text{C}_4$  alkyl;  $\text{C}_1$ - $\text{C}_4$  alkoxy;  $\text{C}_1$ - $\text{C}_4$  hydroxyalkyl;  $\text{C}_1$ - $\text{C}_4$  methoxyalkyl;  $\text{C}_1$ - $\text{C}_4$  monohaloalkyl;  $\text{C}_1$ - $\text{C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ; or  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;

wherein each  $R_5$  is independently H;  $\text{C}_1$ - $\text{C}_3$  alkyl;  $\text{C}_1$ - $\text{C}_3$  monohaloalkyl; or  $\text{C}_1$ - $\text{C}_3$  polyhaloalkyl;

wherein  $\text{L}'$  is  $-\text{NR}_1-\text{L}-$  or

- 37 -



5

wherein L is C<sub>3</sub>-C<sub>9</sub> alkyl; C<sub>3</sub>-C<sub>9</sub> alkenyl; C<sub>3</sub>-C<sub>9</sub> alkynyl;

10




15




20

25



or



15 wherein  $R_1$  is H; or  $C_1$ - $C_3$  straight chained alkyl;

wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

20 wherein one dashed line is a double bond and the other dashed line is a single bond;

25 wherein each  $R_6$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_5$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_6$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

30 wherein  $i$  is an integer from 1 to 4; wherein  $n$  is an integer from 0 to 3; wherein  $m$  is an integer from 0 to 3;

wherein  $K$  is  $-CH_2-NR_{10}-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-(CH_2)_3-$ ;  $-CH_2-NH-CO-NH-(CH_2)_3-$ ;  $-CO-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-(CH_2)_3-$ ;  $-CH_2-NH-CS-NH-(CH_2)_3-$ ;  $-CS-NH-$



-39-


$\text{CHR}_7 - (\text{CH}_2)_j -$ ; or  $-\text{CH}_2 - \text{NR}_{10} - \text{CS} - \text{CHR}_7 - (\text{CH}_2)_j$ ; or  $-\text{CH}_2 - \text{N} = \text{CSR}_1 - \text{NH} - (\text{CH}_2)_j$ ;


wherein  $j$  is an integer from 0 to 3;

5

wherein  $R_7$  is H;  $\text{C}_1 - \text{C}_6$  alkyl;  $\text{CH}_2\text{OR}_5$ ;  $(\text{CH}_2)_p\text{NHCO}_2\text{R}_5$ ;  $(\text{CH}_2)_p\text{NHSO}_2\text{R}_5$ ;  $\text{CH}_2\text{N}(\text{R}_{11})_2$ ; phenyl; pyridyl; thiophenyl; or naphthyl;

wherein W is





wherein Q is O; S; N; NR<sub>9</sub>; or C(R<sub>5</sub>)<sub>2</sub>;

5 wherein b is an integer from 1 to 2;

wherein R<sub>8</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; =O;  
 C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub>  
 methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl;  
 10 N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>;  
 N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;

wherein R<sub>9</sub> is H; C<sub>1</sub>-C<sub>3</sub> alkyl; COR<sub>5</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>;


15 wherein R<sub>10</sub> is H; or C<sub>1</sub>-C<sub>6</sub> alkyl;

wherein R<sub>11</sub> is H; COR<sub>5</sub>; COR<sub>12</sub>; SO<sub>2</sub>R<sub>5</sub>; SO<sub>2</sub>R<sub>12</sub>; and

wherein  $R_{12}$  is phenoxy; phenyl, pyridyl; thiophenyl; or naphthyl; wherein the phenoxy, phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$  alkylthio,  $NO_2$ , phenyl, pyridyl or thiophenyl; or a pharmaceutically acceptable salt thereof.


In an embodiment of the present invention the feeding disorder may be obesity or bulimia. In another embodiment of the present invention the subject is a vertebrate, a mammal, a human or a canine. The invention also provides for the decrease in the consumption of food by the subject by the compound inhibiting the activity of the subject's Y5 receptor.

The invention further provides a method of treating a feeding disorder in a subject which comprises administering to the subject an amount of one of the following compounds:




-43-

5




10




15

20




25

30




35


5



10




15



20


25




30

35

5




10




15

20



25 or

30



This invention also provides a method for treating a disorder in a subject which is alleviated by administering to the subject an amount of a compound described herein which is a Y5 receptor antagonist.

This invention additionally provides a method of treating obesity in a subject which comprises administering to the subject an amount of a Y5 receptor antagonist compound described herein.

5

This invention additionally provides a method of treating non-feeding disorders in a subject which comprises administering to the subject an amount of a compound described herein which is a Y5 receptor antagonist.

10

This invention further provides that any of the methods for treating may comprise administering to the subject a plurality of compounds described herein.

15

The invention also provides for the (-) and (+) enantiomers of the compounds of the subject application described herein. Included in this invention are

pharmaceutically acceptable salts and complexes of all of the compounds described herein. The salts include but are

20

not limited to the acids and bases listed herein. The following inorganic acids; hydrochloric acid, hydrofluoric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and boric acid. The organic acids; acetic acid,

25

trifluoroacetic acid, formic acid, oxalic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, maleic acid, citric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzoic acid, glycolic acid, lactic acid and mandelic acid. The following

30

inorganic bases; ammonia, hydroxyethylamine and hydrazine. The following organic bases; methylamine, ethylamine, propylamine, dimethylamine, diethylamine, trimethylamine, triethylamine, ethylenediamine, hydroxyethylamine, morpholine, piperazine and guanidine. This invention

35

further provides for the hydrates and polymorphs of all of the compounds described herein.



This invention further provides for the metabolites and precursors of the compounds of the present invention. The in vivo actions of numerous enzymes responsible for the generation of metabolites of pharmaceutical compounds are well-known in the art. For example, ethers may be modified to alcohols, or esters may be modified by esterases to yield acids as products. Knowledge of the activities of endogenous enzymes also allows the design of precursors or prodrugs of the compounds of the present invention, which when administered to a subject, such as a vertebrate or a human, are expected to yield metabolites which include the compounds of the present invention. For example, secondary amines may be modified by various substituents, such as methyl, alkanoyl, aroyl, or alkyl or aryl carbamates may be formed, which are expected to yield the compounds of the present invention when acted upon in vivo by endogenous enzymes. Such modifications are intended only as illustrative examples, and are not intended to limit the scope of the present invention, as such modifications and techniques therefor are well-known in the art.

The invention also provides a pharmaceutical composition comprising a therapeutically effective amount of the compounds described above and a pharmaceutically acceptable carrier. In the subject invention a "therapeutically effective amount" is any amount of a compound which, when administered to a subject suffering from a disease against which the compounds are effective, causes reduction, remission, or regression of the disease. In one embodiment the therapeutically effective amount is an amount from about 0.01 mg per subject per day to about 500 mg per subject per day, preferably from about 0.1 mg per subject per day to about 60 mg per subject per day and most preferably from about 1 mg per subject per day to

about 20 mg per subject per day. In the practice of this invention the "pharmaceutically acceptable carrier" is any physiological carrier known to those of ordinary skill in the art useful in formulating pharmaceutical compositions.

5

In another embodiment the pharmaceutical carrier may be a liquid and the pharmaceutical composition would be in the form of a solution. In yet another embodiment, the pharmaceutically acceptable carrier is a solid and the composition is in the form of a powder or tablet. In a further embodiment, the pharmaceutical carrier is a gel and the composition is in the form of a suppository or cream. In a further embodiment the compound may be formulated as a part of a pharmaceutically acceptable transdermal patch.

10

15

A solid carrier can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

20

25

30

35

Liquid carriers are used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized

compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are useful in sterile liquid form compositions for parenteral administration. Sterile liquid carriers can also be utilized for intranasal administration, for example with the use of a pressurized composition, or for inhalatory administration. The liquid carrier for pressurized compositions can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by for example, intramuscular, intrathecal, epidural, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. The compounds may be prepared as a sterile solid composition which may be dissolved or suspended at the time of administration using sterile water, saline, or other appropriate sterile injectable medium. Carriers are intended to include necessary and

-50-

inert binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings.

5 The compound can be administered orally in the form of a sterile solution or suspension containing other solutes or suspending agents, for example, enough saline or glucose to make the solution isotonic, bile salts, acacia, gelatin, sorbitan monoleate, polysorbate 80 (oleate esters of sorbitol and its anhydrides copolymerized with ethylene  
10 oxide) and the like.

The compound can also be administered orally either in liquid or solid composition form. Compositions suitable for oral administration include solid forms, such as  
15 pills, capsules, granules, tablets, and powders, and liquid forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions, and suspensions.

20 Optimal dosages to be administered may be determined by those skilled in the art, and will vary with the particular compound in use, the strength of the preparation, the mode of administration, and the advancement of the disease condition. Additional factors  
25 depending on the particular subject being treated will result in a need to adjust dosages, including subject age, weight, gender, diet, and time of administration.

30 One skilled in the art will readily appreciate that appropriate biological assays will be used to determine the therapeutic potential of the claimed compounds for treating the above noted disorders.

-51-


5 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

Experimental DetailsSynthetic Methods


The compounds of the present invention may be synthesized according to the methods described in Schemes 1-4, as described herein. It is generally preferred that the respective product of each process step, as described hereinbelow, is separated and/or isolated prior to its use as starting material for subsequent steps. Separation and isolation can be effect by any suitable purification procedure such as, for example, evaporation, crystallization, column chromatography, thin layer chromatography, distillation, etc. While preferred reactants have been identified herein, it is further contemplated that the present invention would include chemical equivalents to each reactant specifically enumerated in this disclosure.

Temperatures are given in degrees Centigrade (°C). The structure of final products, intermediates and starting materials is confirmed by standard analytical methods, e.g., microanalysis and spectroscopic characteristics (e.g. MS, IR, NMR). Unless otherwise specified, chromatography is carried out using silica gel. Flash chromatography refers to medium pressure column chromatography according to Still et al., J. Org. Chem. 43, 2921 (1978).

## Scheme 1




Scheme 2






- 55 -

Scheme 3

Where K = -CONHCHR<sub>7</sub>(CH<sub>2</sub>)<sub>j</sub>-Where K = -CH<sub>2</sub>NHCHR<sub>7</sub>(CH<sub>2</sub>)<sub>j</sub>-

Scheme 4



Synthesis of Compounds According to Scheme 1

Preparation of the compounds of the present invention having the structure shown in Formula 1-2, Scheme 1, was carried out using well-known methodology for the preparation of a sulfonamide or sulfamide from an amine. Preferably the appropriate arylsulfonyl or arylsulfamoyl halide, preferably the chloride (i.e., Ar-X-SO<sub>2</sub>Cl), is reacted with a monoprotected linear or cyclic alkylamine (Krapcho and Kuell, Synth. Comm. 20(16):2559-2564, 1990) comprising H<sub>2</sub>N-L-K'', where K'' comprises methylene, in the presence of a base such as a tertiary amine, e.g., triethylamine, dimethylaminopyridine, pyridine or the like, in an appropriate solvent (e.g. CHCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>) as shown in Scheme 1, step A, followed by deprotection of the resulting amine as shown in Scheme 1, Step B, all under mild conditions (typically room temperature), to yield the deprotected amine of Formula 1-1. Alternatively, the primary amine H<sub>2</sub>N-L may be replaced with a secondary amine wherein L comprises a piperidine. The arylsulfonyl or arylsulfamoyl halides are either known in the art or can be prepared according to methods well known in the art.

The deprotected amine may be converted to the product amine of Formula 1-2 by either a single step or two step reductive amination with an aryl substituted aldehyde W-CHO as shown in Scheme 1, Step C, in the presence of a solvent such as toluene or dioxane, at elevated temperature, followed by reduction using sodium borohydride in a solvent such as ethanol. The K'' amine and the aldehyde carbon attached to W together form K in the product.

Compounds of Formula 1-3 in Scheme 1, wherein R<sub>i</sub> is H and j=0 and K comprises an amide, may be synthesized from the compound of Formula 1-1 by amidation using suitable

methods such as those taught in "The Peptides," Vol. 1 (Gross and Meinehofer, Eds. Acaemic Press, N.Y., 1979). For example, the compound of Formula 1-1 may be treated with a carboxylic acid derivative of W in the presence of  
5 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) and dimethylaminopyridine (DMAP) in a suitable solvent such as  $\text{CH}_2\text{Cl}_2$  as shown in Scheme 1, Step D, at room temperature in an inert atmosphere of argon or nitrogen, to yield the amide compound of Formula 1-3. As  
10 in the previous method, the K' amine and the carboxylic acid carbon attached to W together form K in the product.

Alternatively, the compound of Formula 1-3 may be synthesized by acylation of the amine of Formula 1-1 using  
15 the acid chloride of W, i.e.,  $\text{WCOCl}$  or  $\text{W}(\text{CH}_2)_j\text{COCl}$  where j is an integer from 1 to 3, in a solvent such as  $\text{CH}_2\text{Cl}_2$  and a suitable tertiary amine such as triethylamine, at room temperature. Again, the K' amine and the acid  
20 chloride carbon attached to W together form K in the product.

This method also provides an alternative path to the compounds of Formula 1-2 by reduction of the amide of  
25 Formula 1-3 using borane-tetrahydrofuran (THF) complex, in THF as shown in Scheme 1, Step E, at elevated temperature in an inert atmosphere.

Compounds of Formula 1-4 in Scheme 1 where K comprises a ureido moiety may be synthesized by urea formation between  
30 the the compound of Formula 1-1 and a substituted aryl isocyanate or aryl carbamate, as shown in Scheme 1, Step F, in a suitable solvent and a suitable tertiary amine such as triethylamine and N-methylmorpholine, at room  
35 temperature in an inert atmosphere. The ureido moiety comprising K is formed between the K' amine and the

-59-

isocyanate (or isothiocyanate) attached to W, or between K'' and the carbamate derivative of W. Alternatively, compounds containing a thiourea moiety instead of a urea moiety may be synthesized similarly by simply replacing the aryl isocyanate described above with an aryl isothiocyante.

Suitable aryl carbamates may be of the form  $WCO_2Ar'$  where W is for example



and  $Ar'$  may be for example 4-nitrophenyl.

#### Synthesis of the compounds of Table 1

As an illustrative example of the synthesis of the compounds shown in Table 1, the synthesis of Example 1 from Table 1, is provided below:

**N-[6{(Naphthalen-2-ylmethyl)-amino}-hexyl]-2-nitrobenzenesulfonamide**

#### 25 Step A, Scheme 1

**[6-(2-Nitrobenzenesulfonylamino)-hexyl]-carbamic acid t-butyl ester:**

To a stirred solution of N-Boc 1, 6-diaminohexane hydrochloride (1.51 g, 6 mmol) and triethyl amine (1.31 g, 13 mmol) in 50 mL methylene chloride was added 2-Nitrobenzenesulfonyl chloride (1.326 g, 6 mmol). The reaction mixture was stirred for 6 h at room temperature, quenched with brine, and extracted with methylene chloride (2x50 mL). The organic layer was washed with brine (a saturated solution of sodium chloride in water, unless

-60-

otherwise specified), dried over anhydrous sodium sulfate, and concentrated in vacuo to yield the titled compound as yellow oil (2.1 g, 87%).

5      **Step B, Scheme 1**

**N-(6-Aminohexyl)-2-nitrobenzenesulfonamideHydrochloride:**

To a stirred solution of [6-(2-Nitrobenzenesulfonylamino)-hexyl]-carbamic acid t-butyl ester (2.0 g, 4.9 mmol) in 25 mL of methylene chloride at room temperature was added 3  
10      mL of saturated HCl solution in ethyl acetate and stirred for 4 h. The precipitated solid was filtered to yield the titled compound as white solid (1.58 g, 95%); mp 161-162 °C.

15      **Step C, Scheme 1**

**N-[6{(Naphthalen-2-ylmethyl)-amino}-hexyl]-2-nitrobenzenesulfonamide:**

A mixture of N-(6-Aminohexyl)-2-nitrobenzenesulfonamide hydrochloride (0.67 g, 2.0 mmol) and 2-naphthaldehyde (0.32  
20      g, 2.1 mmol) in 75 mL of toluene was refluxed using a Dean Stark trap for 20 h. The reaction mixture was concentrated in vacuo. The residue was dissolved in ethanol (40 mL) and sodium borohydride (0.020 g, 6.0 mmol) was added. After the reaction was complete (6 h), the solvent was evaporated  
25      and residue taken up in 30 mL of saturated sodium chloride solution and extracted with ethyl acetate (3x40mL), washed with saturated sodium chloride solution, dried over sodium sulfate and concentrated to afford an oil. The oil was flash chromatographed over silica gel to afford the titled  
30      compound (0.16 g, 37%) which was converted into hydrochloride salt (mp 169°C).

An example of the use of the alternative path to the compounds of Formula 1-2 by reduction of the amide of  
35      Formula 1-3 using borane-THF complex as previously

described for Scheme 1, Step E, is provided for Example 15 of Table 1, as follows:

**N-[6{(1, 2, 3, 4-tetrahydronaphthalen-2-ylmethyl)-amino}-hexyl]-2-aminobenzenesulfonamide**

To a solution of 1, 2, 3, 4-tetrahydro-2-naphthalencarboxylic acid[6-(2-nitrobenzenesulfonylamino)-hexyl]-amide (0.090 g, 0.19 mmol) in tetrahydrofuran (5 mL) cooled to 0 °C was added 2 mL 1M solution of borane:THF complex and the reaction mixture was refluxed for 12h. The reaction mixture was cooled in an ice bath and quenched with 2 mL of 1N HCl. The reaction mixture was neutralized with 10% aqueous sodium hydroxide solution and extracted with ethyl acetate (3x25 mL). The organic phase was washed with brine, dried over sodium sulfate, and evaporated in vacuo to afford an oil which was purified by preparative TLC to afford the titled compound (0.06 g, 70%); hydrochloride salt mp (162-163 °C).

Using appropriately substituted starting materials, the other Examples shown in Table 1 were synthesized as described above, with the exception of Example 52. The compound of Example 52 in Table 1 was synthesized similarly, except that before deprotection of the amine of Formula 1-1 in Scheme 1 Step B, the sulfonyl nitrogen was alkylated with methyl iodide in dimethylformamide at room temperature to afford the N-methylated sulfonamide product (Sato et al., 1995), which was subsequently deprotected as in Scheme 1, Step B, for use in the remainder of Scheme 1. Other n-alkyl derivatives may be prepared similarly, using an n-alkyl halide in an inert solvent such as dimethylformamide as described above.

Table 1





















| No. | Ar  | X | R <sub>1</sub> | L                                  | K                                 | W   | mp     | Analysis  |
|-----|---|---|----------------|------------------------------------|-----------------------------------|---|--------|---|
| 1   |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 169    | C <sub>23</sub> H <sub>26</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ HCl + 0.05 CHCl <sub>3</sub> |
| 2   |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 174    | C <sub>27</sub> H <sub>30</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                          |
| 3   |  | - | H              | -(CH <sub>2</sub> ) <sub>4</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 194-5  | C <sub>28</sub> H <sub>28</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ HCl + 0.04 CHCl <sub>3</sub> |
| 4   |   | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 190-1  | C <sub>27</sub> H <sub>30</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                          |
| 5   |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 149-50 |   |
| 6   |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 158    | C <sub>24</sub> H <sub>22</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl            |
| 7   |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 111-12 |   |
| 8   |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 139    | C <sub>21</sub> H <sub>20</sub> N <sub>2</sub> O <sub>6</sub> S<br>+ HCl + 0.1H <sub>2</sub> O    |
| 9   |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 172-4  | C <sub>24</sub> H <sub>27</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl            |
| 10  |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 167    | C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl            |



Table 1 (cont)





















| No. | Ar  | X | R <sub>1</sub> | L                                  | K                                 | W   | mp     | Analysis   |
|-----|---|---|----------------|------------------------------------|-----------------------------------|---|--------|--|
| 11  |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 139-40 | C <sub>27</sub> H <sub>31</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                                       |
| 12  |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 194-6  | C <sub>23</sub> H <sub>28</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                                       |
| 13  |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 119-20 | C <sub>25</sub> H <sub>30</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ HCl                                       |
| 14  |  | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |  | 120-1  | C <sub>24</sub> H <sub>30</sub> N <sub>2</sub> O <sub>3</sub> S<br>+ HCl                                       |
| 15  |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 162-63 | C <sub>23</sub> H <sub>28</sub> N <sub>3</sub> O <sub>2</sub> S<br>+ 2.0 HCl                                   |
| 16  |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 141    | C <sub>24</sub> H <sub>30</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl + 0.1 CH <sub>2</sub> Cl <sub>2</sub> |
| 17  |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 183-6  | C <sub>23</sub> H <sub>27</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl                                       |
| 18  |    | - | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 181-4  | C <sub>26</sub> H <sub>31</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl                                       |
| 19  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 158    | C <sub>29</sub> H <sub>34</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                                       |
| 20  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |    | 145-6  | C <sub>29</sub> H <sub>34</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                                       |

Table I (cont)

| No. | Ar | X | R <sub>1</sub> | L                                  | K                                 | W | mp     | Analysis  |
|-----|----|---|----------------|------------------------------------|-----------------------------------|---|--------|---|
| 21  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 117-20 | C <sub>26</sub> H <sub>30</sub> N <sub>3</sub> O <sub>4</sub> SF <sub>3</sub><br>+ HCl                          |
| 22  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 153-5  | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl                          |
| 23  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 166-8  | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>2</sub> S<br>+ 2.0 HCl                                    |
| 24  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 120-2  | C <sub>26</sub> H <sub>34</sub> N <sub>2</sub> O <sub>3</sub> S<br>+ HCl + 0.05 CHCl <sub>3</sub>               |
| 25  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 134-6  | C <sub>25</sub> H <sub>35</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl  |
| 26  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 129-31 | C <sub>25</sub> H <sub>37</sub> N <sub>3</sub> O <sub>2</sub> S<br>+ 2.0 HCl                                    |
| 27  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 113-4  | C <sub>26</sub> H <sub>38</sub> N <sub>3</sub> O <sub>3</sub> S<br>+ HCl + 0.1 CHCl <sub>3</sub>                |
| 28  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 250-1  | C <sub>26</sub> H <sub>32</sub> N <sub>3</sub> O <sub>2</sub> S<br>+ HCl  |
| 29  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 195    | C <sub>29</sub> H <sub>41</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl + 0.25 H <sub>2</sub> O                |
| 30  |    | - | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCH <sub>2</sub> |   | 172    | C <sub>29</sub> H <sub>32</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl + 0.25 CH <sub>2</sub> Cl <sub>2</sub> |

Table 1(cont)































| No. | Ar  | X | R <sub>1</sub> | L   | K                                 | W   | mp          | Analysis   |
|-----|---|---|----------------|---|-----------------------------------|---|-------------|--|
| 31  |  | - | H              |  | CH <sub>2</sub> NHCH <sub>2</sub> |  | 210         | C <sub>26</sub> H <sub>36</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl   |
| 32  |  | - | H              |  | CH <sub>2</sub> NHCH <sub>2</sub> |  | 220         | C <sub>26</sub> H <sub>36</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl + 0.15 CH <sub>2</sub> Cl <sub>2</sub>  |
| 33  |  | - | H              |  | CH <sub>2</sub> NHCH <sub>2</sub> |  | 201         | C <sub>26</sub> H <sub>32</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl + 0.3 CHCl <sub>3</sub>                 |
| 34  |    | - | H              |   | CH <sub>2</sub> NHCH <sub>2</sub> |   | 194-5       | C <sub>25</sub> H <sub>26</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl + 0.1 CHCl <sub>3</sub>                 |
| 35  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | Hygroscopic | C <sub>25</sub> H <sub>26</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl + 0.2 C <sub>6</sub> H <sub>14</sub>    |
| 36  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | 200-2       | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ HCl   |
| 37  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | 216-7       | C <sub>26</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl                           |
| 38  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | 171-4       | C <sub>26</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl + 0.075 CHCl <sub>3</sub> |
| 39  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | 75-8        | C <sub>26</sub> H <sub>33</sub> N <sub>2</sub> O <sub>2</sub> SF <sub>3</sub><br>+ HCl + 0.05 CHCl <sub>3</sub>  |
| 40  |    | - | H              |    | CH <sub>2</sub> NHCH <sub>2</sub> |    | 175-7       | C <sub>26</sub> H <sub>35</sub> N <sub>3</sub> O <sub>2</sub> S<br>+ 2 HCl + 0.8 Et <sub>2</sub> O               |

Table 1(cont)





































| No. | Ar  | X | R <sub>1</sub> | L   | K                          | V   | mp          | Analysis   |
|-----|---|---|----------------|---|----------------------------|---|-------------|--|
| 41  |  | - | H              |  | $\text{CH}_2\text{NHCH}_2$ |  | Hygroscopic | $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_2\text{S}$<br>+ HCl                          |
| 42  |  | - | H              |  | $\text{CH}_2\text{NHCH}_2$ |  | 118-20      | $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_2\text{S}$<br>+ 2.6 HCl                      |
| 43  |   | - | H              |   | $\text{CH}_2\text{NHCH}_2$ |  | 115 dec     | $\text{C}_{23}\text{H}_{31}\text{N}_6\text{O}_4\text{S}$<br>+ 2.4 HCl                      |
| 44  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 89 dec      | $\text{C}_{24}\text{H}_{28}\text{N}_4\text{O}_4\text{S}$<br>+ 2 HCl                        |
| 45  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 104-6       | $\text{C}_{24}\text{H}_{28}\text{N}_4\text{O}_4\text{S}$<br>+ 2 HCl + 0.25 $\text{CHCl}_3$ |
| 46  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 78-80       | $\text{C}_{24}\text{H}_{32}\text{N}_4\text{O}_4\text{S}$<br>+ 2 HCl + 0.65 $\text{CHCl}_3$ |
| 47  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 249-51      | $\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}_6\text{S}$<br>+ 1 HCl + 0.1 $\text{CHCl}_3$  |
| 48  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 220-21      | $\text{C}_{22}\text{H}_{27}\text{N}_3\text{O}_6\text{S}$<br>+ HCl + 0.05 $\text{CHCl}_3$   |
| 49  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 62-5        | $\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}_6\text{S}$<br>+ HCl + 0.5 $\text{CHCl}_3$    |
| 50  |    | - | H              |    | $\text{CH}_2\text{NHCH}_2$ |    | 196-7       | $\text{C}_{24}\text{H}_{31}\text{N}_3\text{O}_5\text{S}$<br>+ HCl                          |

Table 1 (cont)

| No. | Ar  | X | R <sub>1</sub>  | L   | K                                 | W   | m.p   | Analysis  |
|-----|---|---|-----------------|---|-----------------------------------|---|-------|---|
| 51  |  | - | H               |  | CH <sub>2</sub> NHCH <sub>2</sub> |  | 57    | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>5</sub> S<br>+ HCl + 0.13 CHCl <sub>3</sub> |
| 52  |  | - | CH <sub>3</sub> |  | CH <sub>2</sub> NHCH <sub>2</sub> |  | 235-6 | C <sub>30</sub> H <sub>34</sub> N <sub>2</sub> O <sub>2</sub> S<br>+ HCl                          |

### Synthesis of the compounds of Table 2

As an illustrative example of the synthesis of the compounds shown in Table 2 as shown in Scheme 1, Step D, the synthesis of Example 53 from Table 2, is provided below:

1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-(2-nitrobenzenesulfonylamino)-hexyl]-amide

#### Step D, Scheme 1

1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-(2-nitrobenzenesulfonylamino)-hexyl]-amide:

A mixture of N-(6-aminohexyl)-2-nitrobenzenesulfonamide hydrochloride (0.2 g, 0.58 mmol), EDC (0.252 g, 1.31 mmol), and DMAP (0.14 g, 1.21 mmol) in methylene chloride (30 mL) was stirred at room temperature for 0.5h. 1, 2, 3, 4-tetrahydronaphthalen-2-carboxylic acid (0.114 g, 0.65 mmol) was added to the reaction mixture and stirred at room temperature until the completion of the reaction (determined by TLC). The reaction mixture was washed with saturated ammonium chloride (3x30 mL), dried over sodium sulfate and concentrated in vacuo. The residue was flash chromatographed over silica gel to afford an oil (0.25 g, 93%).

Other compounds of Formula 1-3, as shown in Table 2, were synthesized using the methods described above, except for Examples 59 and 63. Example 59 was synthesized as shown in Scheme 1, except for the use of sulfamyl chloride starting material Ar-NH-SO<sub>2</sub>Cl instead of the sulfonyl chloride Ar-SO<sub>2</sub>Cl (Benson and Spillane, Chem. Rev. 80:151-186, 1980).

Example 63 of Table 2 was synthesized from the compound of Example 53 of Table 2 by reduction of the aryl nitro moiety using tin chloride in HCl/Acetic acid/H<sub>2</sub>O at room

temperature, to afford the amino aryl derivative (such as in Example 56, Table 2), which was then sulfonylated with alkyl sulfonyl chloride to afford the bis-sulfonylated compound of Example 63 as follows:

5

1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-{2-(bismethanesulfonylamino)benzenesulfonylamino}]-hexyl]-amide

10

(a) 1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-(2-aminobenzenesulfonylamino)-hexyl]-amide:

15

To a solution of 1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-(2-nitrobenzenesulfonylamino)-hexyl]-amide (0.54 g, 1.17 mmol) in 5 mL of glacial acetic acid stirred at 0 °C was added a solution of tin chloride (1.32 g, 5.88 mmol) in 5 mL conc. HCl and 1 mL of water. The reaction mixture was warmed to room temperature and stirred for 3 h and poured over crushed ice (50 g). The reaction mixture was neutralized with 10% sodium hydroxide and extracted with chloroform (5x50 mL), dried over sodium sulfate and concentrated to afford yellow oil (0.48 g, 95%).

20

25

(b) 1, 2, 3, 4-Tetrahydro-2-naphthalencarboxylic acid[6-{2-(bismethanesulfonylamino)benzenesulfonylamino}]-hexyl]-amide:

30

To a solution of 1, 2, 3, 4-tetrahydro-2-naphthalencarboxylic acid[6-(2-aminobenzenesulfonylamino)-hexyl]-amide (0.075 g, 0.17 mmol) in chloroform (3 mL), and triethylamine (0.1 g, 1 mmol) stirred at 0 °C was added methane sulfonyl chloride (30 mL, 0.35 mmol) and the reaction mixture was stirred at room temperature for 3 h. Solvent was evaporated and purifications by preparative TLC afforded the titled compound as white solid (0.1 g, 97%); mp 70 °C.

35

-70-

The amine of Example 56, Table 2, further may be acylated in the same manner as above in Example 63 but using suitable alkyl acyl chlorides instead of an alkyl sulfonyl chloride.



Table 2

| No. | Ar | X  | R <sub>1</sub> | L                                  | K                    | W | mp   | Analysis  |
|-----|----|----|----------------|------------------------------------|----------------------|---|------|---|
| 53  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>23</sub> H <sub>20</sub> N <sub>3</sub> O <sub>5</sub> S<br>+ 0.35 CH <sub>2</sub> Cl <sub>2</sub>             |
| 54  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>24</sub> H <sub>32</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ 0.6 CHCl <sub>3</sub>                            |
| 55  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>24</sub> H <sub>26</sub> N <sub>3</sub> O <sub>5</sub> SF <sub>3</sub><br>+ 0.2 C <sub>6</sub> H <sub>14</sub> |
| 56  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | 72-4 | C <sub>23</sub> H <sub>31</sub> N <sub>3</sub> O <sub>3</sub> S<br>+ 0.25 CH <sub>2</sub> Cl <sub>2</sub>             |
| 57  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | 132  | C <sub>23</sub> H <sub>28</sub> N <sub>4</sub> O <sub>5</sub> SF <sub>3</sub><br>+ HCl + 0.05 CHCl <sub>3</sub>       |
| 58  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | 98   | C <sub>22</sub> H <sub>28</sub> N <sub>4</sub> O <sub>5</sub> S<br>+ HCl + 0.1 CHCl <sub>3</sub>                      |
| 59  |    | NH | H              | -(CH <sub>2</sub> ) <sub>4</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>22</sub> H <sub>28</sub> N <sub>4</sub> O <sub>5</sub> S<br>+ 0.35 CH <sub>2</sub> Cl <sub>2</sub>             |
| 60  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>23</sub> H <sub>25</sub> N <sub>3</sub> O <sub>5</sub> S   |
| 61  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | 105  | C <sub>23</sub> H <sub>25</sub> N <sub>3</sub> O <sub>5</sub> S   |
| 62  |    | -  | H              | -(CH <sub>2</sub> ) <sub>7</sub> - | CH <sub>2</sub> NHCO |   | Oil  | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>5</sub> S<br>+ 0.15 CHCl <sub>3</sub>                           |
| 63  |    | -  | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO |   | 70   | C <sub>25</sub> H <sub>35</sub> N <sub>3</sub> O <sub>7</sub> S <sub>3</sub>  |

Table 2 (cont)

| No. | Ar | X | R <sub>1</sub> | L | K                    | W | mp    | Analysis   |
|-----|----|---|----------------|---|----------------------|---|-------|--|
| 64  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 194-7 | C <sub>25</sub> H <sub>33</sub> N <sub>3</sub> O <sub>3</sub> S<br>+ HCl + 0.2 CHCl <sub>3</sub>         |
| 65  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 78-80 | C <sub>25</sub> H <sub>39</sub> N <sub>3</sub> O <sub>6</sub> S<br>+ HCl + 0.3 CHCl <sub>3</sub>         |
| 66  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 86-87 | C <sub>29</sub> H <sub>32</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ HCl + 0.35 CHCl <sub>3</sub>        |
| 67  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 59-61 | C <sub>29</sub> H <sub>32</sub> N <sub>2</sub> O <sub>4</sub> S<br>+ 0.2 CHCl <sub>3</sub>               |
| 68  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 150-3 | C <sub>23</sub> H <sub>25</sub> N <sub>5</sub> O <sub>5</sub> S<br>+ 0.25 CHCl <sub>3</sub>              |
| 69  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 62-4  | C <sub>24</sub> H <sub>26</sub> N <sub>4</sub> O <sub>5</sub> S<br>+ 0.3 CHCl <sub>3</sub>               |
| 70  |    | - | H              |   | CH <sub>2</sub> NHCO |   | 85-7  | C <sub>27</sub> H <sub>37</sub> N <sub>3</sub> O <sub>7</sub> S <sub>3</sub><br>+ 0.25 CHCl <sub>3</sub> |

**Synthesis of the compounds of Table 3**

As an illustrative example of the synthesis of the compounds shown in Table 3 and Table 3a, as shown in Scheme 1, Step F, the synthesis of Example 71 from Table 3, is provided below:

**Synthesis of Naphthalene-1-sulfonic acid{6-[3-(1-naphthyl)-uriedo]-hexyl}-amide****Step F, Scheme 1**

To a solution of N-(6-aminohexyl)-1-naphthalenesulfonamide hydrochloride(0.1 g, 0.29 mmol) in 3 mL dimethylformamide and 50 mL N-methyl morpholine was added 1-naphthalene isocyanate(68  $\mu$ L, 0.4 mmol). The reaction mixture was stirred at room temperature for 6 h. Solvent was evaporated in vacuo and residue was purified by preparative TLC to afford white solid(110 mg, 79%);mp 105-106 °C.

An example of the synthesis of a compound of Formula 1-4 in Scheme 1, using an aryl carbamate as previously described, is provided for Example 74 of Table 3, as follows:

**3, 4-Dihydroquinoline-1-carboxylic acid[6-(2-trifluoromethylbenzenesulfonylamino)-hexyl]amide****(a) 3, 4-Dihydro-2H-quinoline-1-carboxylic acid 4-nitrophenyl ester**

To a solution of tetrahydroisoquinoline (3.99 g, 30 mmol) in triethyl amine(6 g, 60 mmol) and dichloromethane(150 mL) cooled in ice bath was added p-nitrophenyl chloroformate (6.06 g, 30 mmol) drop wise over a period of 1h. The reaction mixture was stirred at room temperature for 3h and then washed with water (3x100 mL), dried over sodium sulfate and concentrated to afford an oil which was

trituated with ether:hexane to afford the titled compound as yellow solid (5.9 g, 65%).

5 (b) 3, 4-Dihydroquinoline-1-carboxylic acid[6-(2-trifluoromethylbenzenesulfonylamino)-hexyl]amide:

To a solution of N-(6-Aminohexyl)-2-trifluoromethylbenzene sulfonamide Hydrochloride (0.1 g, 0.27 mmol) dimethyl formamide (3mL) and triethylamine (0.1 g, 1mmol) was added 3, 4-Dihydro-2H-quinoline-1-carboxylic acid 4-nitrophenyl ester (0.09 g, 0.3 mmol) and the reaction mixture was stirred at room temperature for 4h. Solvent was evaporated and purification by preparative TLC afford the titled compound as an oil (0.08g, 61 %).

15 **Synthesis of the compounds of Table 3a**

As illustrative examples of the synthesis of the compounds of Table 3a, as shown in Scheme 1, Step F, the syntheses of Examples 80-82 and 87 is provided below:

20 **Example 80**

1-[1-(Naphthalene-1-sulfonyl)-piperidine-4-ylmethyl]-3-naphthalene-1-ylmethylthiourea

1-Naphthalene-1-ylmethyl-3-piperidine-4-ylmethylthiourea

25 To a solution of 1-naphthalenemethylisothiocyanate (2.8 g, 13.6 mmol) in 100 ml of MeOH-THF solution (1:1 mixture) was added 4-aminomethylpiperidine (3.1 ml, 27.0 mmol) in a portion and the resulting solution was stirred at reflux for 12 h. The reaction mixture was concentrated in vacuo, yielding oily mixture which was subjected to column chromatography (10% MeOH/CHCl<sub>3</sub>) to yield 2.3 g (48%) of the desired product as an oil.

35 1-[1-(Naphthalene-1-sulfonyl)-piperidine-4-ylmethyl]-3-naphthalene-1-ylmethylthiourea

- 75 -

To a solution of the amine (0.34 g, 1.1 mmol) in 10 ml of pyridine was added 1-naphthalenesulfonyl chloride (0.30 ml, 1.3 mmol) in a portion and the resulting mixture was stirred at 25°C for 12 h. The reaction mixture was subjected to column chromatography (50% Hexane/EtOAc) to yield 0.21 g (38%) of the desired product as yellow solid, which was recrystallized from EtOAc to provide 0.15 g of the product as white solid: mp 83 - 85 °C; Anal. Cal. For  $C_{28}H_{29}N_3O_2S_2$  requires C, 66.77; H, 5.80; N, 8.34. Found: C, 64.32; H, 5.89; N, 8.27.

**Example 81****2-methyl-1-[1-(naphthalene-1-sulfonyl)piperidine-4-ylmethyl]-3-naphthalene-1-ylmethylisothiourea**

To a solution of the thiourea of Example 80 (0.11 g, 0.32 mmol) in MeOH was added MeI (0.5 ml, 8.1 mmol) in a portion and the resulting mixture was stirred at 25°C for 12 h. The reaction mixture was concentrated in vacuo, yielding a solid which was recrystallized from EtOH to yield 0.11 g (53%) of the desired product as white solid: mp 111 - 113°C; Anal. Cal. For  $C_{29}H_{31}N_3O_2S_2 \cdot 1.0 HI$  requires C, 54.04; H, 4.89; N, 6.52. Found: C, 52.97; H, 5.01; N, 6.47.

**Example 82****1-[1-(Naphthalene-1-sulfonyl)-piperidine-4-ylmethyl]-3-naphthalene-1-ylmethylurea****1-Naphthalene-1-ylmethyl-3-piperidine-4-ylmethylurea**

To a solution of 1-naphthalenemethylcyanate (1.1 g, 6.0 mmol) in 50 ml of  $CHCl_3$  was added 4-aminomethylpiperidine (0.9 ml, 7.8 mmol) in a portion and the resulting solution was stirred at reflux for 12 h. The reaction mixture was concentrated in vacuo, yielding oily mixture which was purified on column chromatography (5% MeOH/ $CHCl_3$ ) to yield

-76-

the desired product 1.6 g (89%) of the desired product as an oil.

**1-[1-(Naphthalene-1-sulfonyl)-piperidine-4-ylmethyl]-3-naphthalene-1-ylmethylurea**

To a solution of the amine (0.60 g, 2.0 mmol) in 30 ml of pyridine was added 1-naphthalenesulfonyl chloride (0.60 ml, 2.7 mmol) in a portion and the resulting mixture was stirred at 25°C for 12h. The reaction mixture was subjected to column chromatography (CHCl<sub>3</sub>, neat) to yield 0.63 g (65%) of the desired product as light yellow solid, which was recrystallized from EtOAc to provide 0.37 g of the product as white solid: mp 103 - 104°C; Anal. Cal. For C<sub>28</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>S requires C, 64.97; H, 5.99; N, 8.62. Found: C, 66.03; H, 5.83; N, 8.49.

**Example 87**

**1-Naphthalene-1-ylmethyl-3-[1-(2-trifluoromethyl-benzenesulfonyl)-pyrrolidin-3-yl]-urea**

**1-(Naphthalene-1-sulfonyl)-pyrrolidin-3-ylamine**

To a solution of 1-naphthalenesulfonyl chloride (1.0 g, 5.4 mmol) in 50 ml of CH<sub>2</sub>Cl<sub>2</sub> with 2 ml of trimethylamine was added 3-t-butoxycarbonylpyrrolidine (1.5 g, 6.5 mmol) in a portion and the resulting solution was stirred at reflux for 48 h. Reaction mixture was concentrated in vacuo, yielding oily mixture which was partitioned between 100 ml of EtOAc and NaHCO<sub>3</sub> sat'd aqueous solution. Organic layer was separated, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated in vacuo to provide an oil, which was redissolved in 20 ml of CH<sub>2</sub>Cl<sub>2</sub>. Toward this solution was added 1 ml of trifluoroacetic acid and the resulting solution was stirred for 2 h at 25°C. Reaction mixture was concentrated in vacuo, providing a brown oil, which was dissolved in EtOAc and washed with aqueous NaOH.

-77-

Organic layer was concentrated in vacuo to yield an oil which was subjected to column chromatography (30% MeOH/CHCl<sub>3</sub>) to provide 0.17 g (23%) of the desired product.

5

**1-Naphthalene-1-ylmethyl-3-[1-(2-trifluoromethyl-benzenesulfonyl)-pyrrolidin-3-yl]-urea**

The amine (37 mg, 0.12 mmol) and naphthalene-1-ylmethyl-carbamic acid 4-nitro-phenyl ester (61 mg, 0.19 mmol) in 10 5 ml of CH<sub>2</sub>Cl<sub>2</sub> was stirred at 25°C for 12 h. The reaction mixture was subjected to column chromatography (2% MeOH/CHCl<sub>3</sub>) to yield 42 mg (74%) of the desired product as light yellow solid: mp 117 - 119 °C; Anal. Cal. For C<sub>23</sub>H<sub>22</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub>S requires C, 57.85; H, 4.64; N, 8.80. 15 Found: C, 56.16; H, 4.71; N, 8.67.

Additional compounds may be synthesized by substitution of appropriate starting materials, and are not intended to be limited to those compounds shown in Table 3a.

20

Other compounds of Formula 1-4, as shown in Table 3 and Table 3a, were synthesized using the methods described above, using appropriately substituted starting materials.

Table 3













































| No. | Ar  | X | R <sub>1</sub> | L                                  | K                      | W  | mp     | Analysis   |
|-----|---|---|----------------|------------------------------------|------------------------|--|--------|--|
| 71  |  | . | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCONH |  | 105-06 | C <sub>27</sub> H <sub>29</sub> N <sub>3</sub> O <sub>3</sub> S                            |
| 72  |  | . | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCONH |   | 115    | C <sub>23</sub> H <sub>26</sub> N <sub>4</sub> O <sub>6</sub> S                            |
| 73  |  | . | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCONH |   | Oil    | C <sub>24</sub> H <sub>26</sub> N <sub>3</sub> O <sub>4</sub> S<br>+ 0.5 CHCl <sub>3</sub> |
| 74  |  | . | H              | -(CH <sub>2</sub> ) <sub>5</sub> - | CH <sub>2</sub> NHCO   |   | Oil    | C <sub>23</sub> H <sub>26</sub> N <sub>3</sub> O <sub>3</sub> S                            |



Table 3a

| No | Ar  | X | L   | K   | W   | mp      |                            |
|----|---|---|---|---|---|---------|----------------------------|
| 81 |  | - |  |  |  | 111-113 | $C_{27}H_{31}N_3O_2S + HI$ |
| 80 |   | - |   |   |   | 83-85   | $C_{26}H_{29}N_3O_2S_2$    |
| 82 |    | - |    |    |    | 104-106 | $C_{28}H_{33}N_3O_2S$      |
| 83 |    | - |    |    |    | 176-178 | $C_{28}H_{31}N_3O_2S_2$    |
| 84 |    | - |    |    |    | 199-201 | $C_{27}H_{29}N_3O_2S_2$    |
| 85 |    | - |    |    |    | 94-97   | $C_{27}H_{26}F_3N_3O_2S$   |
| 86 |    | - |    |    |    | 88-90   | $C_{27}H_{26}F_3N_3O_2S_2$ |
| 87 |    | - |    |    |    | 117-119 | $C_{27}H_{22}F_3N_3O_2S$   |
| 88 |    | - |    |    |    | 162-164 | $C_{27}H_{27}N_3O_2S$      |

**Synthesis of compounds of Scheme 2**

Compounds in which L is substituted may be produced according to Scheme 2. The carbobenzyloxy-protected amino acid derivative of L may be esterified by thionyl chloride and methanol as shown in Scheme 2, Step A, and the ester subsequently reduced by diisobutylaluminum hydride in THF as shown in Scheme 2, Step B, at -78 °C, to yield alcohol, which is then oxidized by pyridinium chlorochromate in Scheme 2, Step C, in a suitable solvent such as CH<sub>2</sub>Cl<sub>2</sub>, to afford an aldehyde of Formula 2-1. The aldehyde may be treated with ammonia, and then trimethylsilylcyanide (Chakraborty, et al. Tet. Lett. 32(51):7597-7600, 1991) as shown in Scheme 2, Step D, in a suitable solvent such as methanol, to yield the compound of Formula 2-2. The compound of Formula 2-2 may be subjected to sulfonylation as described above in Scheme 1, Step A, to yield the compound of Formula 2-3. The cyano moiety of the compound of Formula 2-3 may be esterified and deblocked to afford compounds of Formula 2-4; or if desired, the compound of Formula 2-3 may be further reduced, or hydrolyzed as appropriate to yield substituted compounds other than those shown in Formula 2-5, which compounds may be further used in any of Steps C, D, E, or F of Scheme 1.

An example of the synthesis of a compound of Formula 2-5 is the synthesis of compound 79:

**7 - [ ( N a p h t h a l e n - 2 - y l m e t h y l ) - a m i n o ] - 2 - ( 2 - n i t r o b e n z e n e s u l f o n y l a m i n o ) - h e p t a n o i c a c i d m e t h y l e s t e r**

**(a) Step A, Scheme 2****6-Benzyloxycarbonylamino-hexanoic Acid Methyl Ester:**

To a solution of 6-benzyloxycarbonylamino-hexanoic acid (10 g, 38 mmol) in methanol (200 mL) at room temperature was added thionyl chloride (11 g, 95 mmol). The reaction

-81-

mixture was then refluxed for 16 h. The solvent was concentrated in vacuo, the residue was dissolved in ethyl acetate (200 mL) and washed with brine (3x150 mL), dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was purified by flash column chromatography (silica, 14% ethyl acetate in hexane) to afford the titled compound (6.2 g, 58%), a colorless liquid.

10 (b) Step B, Scheme 2

(6-Hydroxyhexyl)-carbamic acid benzyl ester:

To a solution of diisobutylaluminum hydride (49 mL, 1.5 M in toluene) in THF (150 mL) cooled to -78 °C was added a solution of 6-benzyloxycarbonylamino-hexanoic acid methyl ester (10 g, 37 mmol) in THF (100 mL). The reaction mixture was stirred for 4 h and methanol (2.5 mL) was added carefully at -78 to -75 °C. After 2 h, the mixture was poured to 250 mL of 1 N HCl solution cooled in ice-bath, extracted with ethyl acetate (300 mL), washed with brine (3x100 mL), dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was purified by flash column chromatography (silica, 50% ethyl acetate in hexane) to yield the titled compound (7.1 g, 80%); white solid, mp 67-68 °C.

25

(c) Step C, Scheme 2

(6-Oxy-hexyl)-carbamic acid benzyl ester:

To a suspension of pyridinium chlorochromate (9.2 g, 43 mmol) and celite (37 g) in methylene chloride (400 mL) was added (6-hydroxyhexyl)-carbamic acid benzyl ester (7.1 g, 24 mmol). The mixture was stirred for 3 h and dry ethyl ether (500 mL) was added to it. The mixture was stirred for additional 0.5 h, filtered through a pad of celite and concentrated in vacuo. The residue was purified by flash column chromatography (silica, 10-25% ethyl acetate in

35

hexane) to yield the titled compound (4.8 g, 79%); colorless light oil.

(d) Step D, Scheme 2

5 (6-Amino-6-cyano-hexyl)-carbamic acid benzyl ester:

Ammonia was bubbled through a stirred solution of (6-oxy-hexyl)-carbamic acid benzyl ester (6.0 g, 24 mmol) in methanol (200 mL) for 2 h and TMSCN (2.9 g, 28 mmol) was added dropwise. The reaction mixture was stirred for 24 h.

10 The solvent was evaporated in vacuo. The residue was purified by flash column chromatography (silica, 90-100% ethyl acetate in hexane) to yield the titled compound (4.3 g, 65%); light yellow oil.

15 (e) (Step A, Scheme 1)

(6-Benzenesulfonylamino-6-cyano-hexyl)-carbamic acid benzyl ester:

Using the general procedure described in step A of scheme 1, (6-amino-6-cyano-hexyl)-carbamic acid benzyl ester (2.8 g, 10 mmol) was sulfonylated with 2-nitrobenzenesulfonyl chloride (2.5 g, 11 mmol) at 0 °C to yield the titled compound (1.7 g, 36%); yellow oil.

(f) Step E, Scheme 2

25 7-Amino-2-nitrobenzenesulfonylamino-heptanoic acid methyl ester:

Dry HCl gas was bubbled to a stirred solution of (6-benzenesulfonylamino-6-cyano-hexyl)-carbamic acid benzyl ester (0.58 g, 1.3 mmol) in dry methanol (50 mL) for 2h.

30 The reaction mixture was cooled and solvent was evaporated in vacuo. Water (40 mL) was added to the reaction mixture and neutralized with 1 N NaOH to pH 9-10, extracted with ethyl acetate (4x100 mL), washed with brine (3x100 mL), dried over anhydrous magnesium sulfate, and  
35 concentrated in vacuo. The residue was purified by

preparative TLC (silica, 10% ammonia (2.0 M in methanol) in chloroform) to yield the titled compound (0.047 g, 10%); yellow thick oil.

- 5        (g) (Steps C, D, E, F, Scheme 1)  
7 - [ (Naphthalen-2-ylmethyl) - amino] - 2 - (2 - nitrobenzenesulfonylamino) - heptanoic acid methyl ester:  
Using the general procedure described in Steps C, D, E, and F of Scheme 1, 7-amino-2-nitrobenzenesulfonylamino-  
10        heptanoic acid methyl ester (0.060 g, 0.17 mmol) was reductively aminated with 2-naphthaldehyde (0.026 g, 0.017 mmol) to afford the titled compound (0.050 g, 60%); yellow oil.
- 15        **Synthesis of compounds according to Scheme 3**  
Other compounds of the present invention may be synthesized according to Scheme 3. After protection of  $H_2N-L-COOH$  with Boc anhydride in  $CH_2Cl_2$ , as shown in Scheme 3, Step A, the protected amine may be amidated with W-K''' as in Scheme 3, Step B, where K''' is an alkylamino ester, using EDC and DMAP in a suitable solvent such as  $CH_2Cl_2$ , to yield compounds of Formula 3-1, where K''' and the carboxylic acid carbonyl of  $H_2N-L-COOH$  together form K. The compounds of Formula 3-1 may be deprotected using well  
20        known methods as shown in Scheme 3, Step C, and further sulfonylated with a sulfonyl halide of Ar, as shown in Scheme 3, Step D, in a suitable solvent such as  $CH_2Cl_2$  and a tertiary amine such as triethylamine, to form the compound of Formula 3-2. The compounds of Formula 3-2 may  
25        be reduced to yield the compounds of Formula 3-3, as shown in Scheme 3, Step E, using borane-tetrahydrofuran (THF) complex, in THF, at elevated temperature in an inert atmosphere.
- 30

A specific example of such a synthesis using Scheme 3 is provided below for Example 75 from Table 4:

5       trans-3-(4-Chloro-phenyl)-2-([4-(naphthalene-1-sulfonylamino)-methyl]-cyclohexanecarbonyl)-amino]-propionic acid methyl ester:

(a) Step A, Scheme 3

trans-4-(tert-Butoxycarbonylamino-methyl)-cyclohexanecarboxylic acid:

10       To a solution of trans-4-(aminomethyl)cyclohexanecarboxylic acid (10 g, 57 mmol) in 1 N NaOH (110 mL) cooled to 0 °C was added a solution of di-tert-butyl dicarbonate (15 g, 69 mmol) in dioxane (50 mL). The reaction mixture was stirred at 0 °C for 12 h.  
15       The reaction mixture was neutralized by 1 N HCl solution to pH 3, extracted with ethyl ether (2x300 mL), washed with brine (2x300 mL), dried over anhydrous magnesium sulfate, and concentrated in vacuo to afford the titled compound (16 g, 100%); white solid, mp 128-9 °C.

20

(b) Step B, Scheme 3

trans-2-{[4-(tert-Butoxycarbonylamino-methyl)-cyclohexanecarbonyl]-amino}3-(4-Chloro-phenyl)-propionic acid methyl ester:

25       Using the general procedure described for the preparation Step D, Scheme 1, trans-4-(tert-butoxycarbonylamino-methyl)-cyclohexanecarboxylic acid (1.1 g, 4.0 mmol) was acylated with D,L-4-chlorophenylalanine methyl ester hydrochloride (1.0 g, 4.0 mmol) to afford the titled  
30       compound (1.9 g, 99%); white solid, mp 178-9 °C.

(c) Step C, Scheme 3

trans-2-[4-(Aminomethyl-cyclohexanecarbonyl)-amino]3-(4-chloro-phenyl)-propionic acid methyl ester hydrochloride:

-85-

Using the general procedure described in step b scheme 1, trans-2-{[4-(tert-butoxycarbonylamino-methyl)-cyclohexanecarbonyl]-amino}3-(4-chloro-phenyl)-propionic acid methyl ester (1.8 g, 4.3 mmol) was deprotected using  
5 HCl in ethyl acetate to afford the titled compound; light yellow solid mp 146-9 °C.

(d) Step D, Scheme 3

trans-3-(4-Chloro-phenyl)-2-({[4-(naphthalene-1-sulfonylamino)-methyl]-cyclohexanecarbonyl}-amino)-propionic acid methyl ester:  
10

Using the general procedure described in example A scheme 1, trans-2-[4-(aminomethyl-cyclohexanecarbonyl)-amino]3-(4-Chloro-phenyl)-propionic acid methyl ester  
15 hydrochloride (0.35 g, 0.86 mmol) was sulfonylated with 1-naphthalenesulfonyl chloride (0.42 g, 91%) to afford the titled ompound; white solid, mp 84-6 °C.

The compound of Example 77, Table 4, was synthesized from  
20 the above compound by borane-THF reduction as follows:













(e) Step E, Scheme 3

Naphthalene-1-sulfonic Acid trans-(4-{[2-(4-Chloro-phenyl)-1-hydroxymethyl-ethylamino]-methyl}-cyclohexylmethyl)-amide:

25 Using the general procedure described in Step H, Scheme 1, trans-3-(4-chloro-phenyl)-2-({[4-(naphthalene-1-sulfonylamino)-methyl]-cyclohexanecarbonyl}-amino)-propionic acid methyl ester (0.30 g, 0.55 mmol) was reduced by borane:THF complex (1.0 M in THF) to afford the  
30 titled compound; colorless oil.

Other compounds of Formula 3-3 or Formula 3-4, as shown in Table 4, where for example, K is substituted with an alcohol or ester, may also be synthesized using the  
35 methods of Scheme 3.

Table 4

| No. | Ar   | X | R <sub>1</sub> | L  | K  | W  | mp    | Analysis   |
|-----|--|---|----------------|--|--|--|-------|--|
| 75  |  | - | H              |  | $\text{PO}_2\text{CH}_3$<br>$\text{CONHCHCH}_2$        |  | 84.6  | $\text{C}_{28}\text{H}_{31}\text{N}_2\text{O}_5\text{SCl}$<br>+ 0.4 $\text{CHCl}_3$  |
| 76  |   | - | H              |   | $\text{PO}_2\text{CH}_3$<br>$\text{CONHCHCH}_2$        |   | Oil   | $\text{C}_{24}\text{H}_{28}\text{N}_3\text{O}_5\text{SCl}$<br>+ 0.15 $\text{CHCl}_3$ |
| 77  |   | - | H              |   | $\text{CH}_2\text{OH}$<br>$\text{CH}_2\text{NHCHCH}_2$ |   | 223-3 | $\text{C}_{27}\text{H}_{33}\text{N}_2\text{O}_5\text{SCl}$<br>+ HCl                  |
| 78  |   | - | H              |   | $\text{CH}_2\text{OH}$<br>$\text{CH}_2\text{NHCHCH}_2$ |   | 263-4 | $\text{C}_{27}\text{H}_{33}\text{N}_2\text{O}_5\text{SCl}$<br>+ HCl                  |



**Synthesis of compounds 89 and 90**

The synthesis of compounds such as 89 and 90 may be accomplished by the method shown in Scheme 4, as described below:

5

**Example 89**

(a) 4-Oxo-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid-(naphthalene-1-ylmethyl)-amide (Scheme 4, product 1):

A mixture of 1-naphthalenemethylamine (1.37 g, 7.2 mmol),  
10 4-oxo-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid  
(1.13 g, 7.2 mmol) ECD (2.87 g, 15 mmol), and DMAP (1.83  
g, 15 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was stirred at room  
temperature. After 12 h, the reaction mixture was  
concentrated and the residue was purified by flash column  
15 chromatography (2-5% MeOH in CH<sub>2</sub>Cl to yield the product  
(2.29 g, 97% light yellow solid, m.p. 155-156°C).

(b) 4-Cyano-4-trimethylsilanyloxy-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid-(naphthalen-1-ylmethyl)-amide (Scheme 4, product 2):

To a mixture of the product of step (a) (2.29 g, 6.95  
mmol) and a catalytic amount of ZnI<sub>2</sub> in dry CH<sub>2</sub>Cl<sub>2</sub> (50 mL)  
at 0°C was added dropwise TMSCN (1.4 g, 1.87 mL, 14 mmol)  
under argon. The reaction mixture was warmed to room  
25 temperature and stirred for 24 h. After concentration of  
the reaction mixture, the residue was purified by flash  
column chromatography (2-5% MeOH in CH<sub>2</sub>Cl to yield the  
product (1.6 g, 54%, 95% after recovering the product from  
(a); light yellow solid, m.p. 55-56°C).

30

(c) 1-Aminomethyl-4-{[(naphthalene-1-ylmethyl)-amino]-methyl}-1,2,3,4-tetrahydronaphthalen-1-ol (Scheme 4, product 3):

To a solution of the product of step (b) (1.4 g, 3.27  
35 mmol) in THF (30 mL) was added dropwise 25 mL of BH<sub>3</sub>-THF

complex (1.0 M in THF) under argon. The reaction mixture was refluxed for 16 h. After cooling to 0°C, 6N HCl was added dropwise to the reaction mixture and the resultant mixture stirred for 24 h at room temperature. After cooling to 0°C, this mixture was neutralized by 1N NaOH to pH 7 to 8, extracted with ethyl acetate (100 mL), washed with water (60 mL x 2), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated. The residue was purified by flash column chromatography (2-5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to yield the product (0.30 g, 27%, light yellow solid).

(d) **N-(1-hydroxy-4-{[(naphthalen-1-ylmethyl)-amino]-methyl}-1,2,3,4-tetrahydronaphthalen-1-ylmethyl-2-nitrobenzenesulfonamide (Scheme 4, product 4):**

To a mixture of the product of step (c) (0.30 g, 0.866 mmol) and Et<sub>3</sub>N (0.35 g, 3.46 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (15 mL) at 0°C was added dropwise a solution of 2-nitrobenzenesulfonyl chloride (0.19 g, 0.866 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The reaction mixture was warmed to room temperature and stirred for 6 h. After concentration of the reaction mixture, the residue was purified by flash column chromatography (2-5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to yield the product (0.43 g, 93%, light yellow solid).

**Example 90 (Scheme 4, product 5)**

A mixture of the compound of Example 89 (0.35 g, 0.658 mmol) and TsOH (0.075 g, 0.395 mmol) in toluene was refluxed for 0.5 h. The reaction mixture was concentrated and purified by TLC chromatography (10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to yield the product (0.217 g, 64%, light yellow solid, m.p. 53-54°C).

**Pharmacological Evaluation of Compounds at Cloned Human Neuropeptide Y-type Receptors.**

The pharmacologic properties of the compounds of the present invention were evaluated at the cloned human neuropeptide Y-type receptors Y1, Y2, Y4, and Y5, or in in vivo studies in rats, using the protocols described below.

5

## MATERIALS AND METHODS

### Cell Culture

COS-7 cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37°C, 5% CO<sub>2</sub>. Stock plates of COS-7 cells were trypsinized and split 1:6 every 3-4 days. Human embryonic kidney 293 cells were grown on 150 mm plates in D-MEM with supplements (minimal essential medium) with Hanks' salts and supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37 °C, 5% CO<sub>2</sub>. Stock plates of 293 cells were trypsinized and split 1:6 every 3-4 days. Mouse fibroblast LM(tk-) cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/mL penicillin/100 µg/mL streptomycin) at 37 °C, 5% CO<sub>2</sub>. Stock plates of LM(tk-) cells were trypsinized and split 1:10 every 3-4 days.

LM(tk-) cells stably transfected with the human Y5 receptor were routinely converted from an adherent monolayer to a viable suspension. Adherent cells were harvested with trypsin at the point of confluence, resuspended in a minimal volume of complete DMEM for a cell count, and further diluted to a concentration of 10<sup>6</sup> cells/ml in suspension media (10% bovine calf serum, 10% 10X Medium 199 (Gibco), 9 mM NaHCO<sub>3</sub>, 25 mM glucose, 2 mM

-90-

5 L-glutamine, 100 units/ml penicillin/100  $\mu$ g/ml streptomycin, and 0.05% methyl cellulose). The cell suspension was maintained in a shaking incubator at 37 °C, 5% CO<sub>2</sub> for 24 hours. Membranes harvested from cells grown in this manner may be stored as large, uniform batches in liquid nitrogen. Alternatively, cells may be returned to adherent cell culture in complete DMEM by distribution into 96-well microtiter plates coated with poly-D-lysine (0.01 mg/ml) followed by incubation at 37 °C, 5% CO<sub>2</sub> for 24 hours. Cells prepared in this manner yielded a robust and reliable NPY-dependent response in cAMP radio-immunoassays as further described hereinbelow.

15 Mouse embryonic fibroblast NIH-3T3 cells were grown on 150 mm plates in Dulbecco's Modified Eagle Medium (DMEM) with supplements (10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100  $\mu$ g/ml streptomycin) at 37 °C, 5% CO<sub>2</sub>. Stock plates of NIH-3T3 cells were trypsinized and split 1:15 every 3-4 days.

20 Sf9 and Sf21 cells were grown in monolayers on 150 mm tissue culture dishes in TMN-FH media supplemented with 10% fetal calf serum, at 27°C, no CO<sub>2</sub>. High Five insect cells were grown on 150 mm tissue culture dishes in Ex-Cell 400™ medium supplemented with L-Glutamine, also at 27°C, no CO<sub>2</sub>.

#### Transient Transfection

30 All receptor subtypes studied (human and rat Y1, human and rat Y2, human and rat Y4, human and rat Y5) were transiently transfected into COS-7 cells by the DEAE-dextran method, using 1  $\mu$ g of DNA /10<sup>6</sup> cells (Cullen, 1987). The human Y1 receptor was prepared using known methods (Larhammar, et al., 1992).

35

Stable Transfection

Human Y1, human Y2, and rat Y5 receptors were co-transfected with a G-418 resistant gene into the human embryonic kidney 293 cell line by a calcium phosphate transfection method (Cullen, 1987). Stably transfected cells were selected with G-418. Human Y4 and human Y5 receptors were similarly transfected into mouse fibroblast LM(tk-) cells and NIH-3T3 cells.

Binding of the compounds of the present invention to the human Y1, Y2, Y4 and Y5 receptors was evaluated using stably transfected 293 or LM(tk-) cells as described above. Stably transfected cell lines which may be used for binding assays include, for example, for the human Y1 receptor, 293-hY1-5 (deposited June 4, 1996, under ATCC Accession No. CRL-12121), for the human Y2 receptor, 293-hY2-10 (deposited January 27, 1994, under ATCC Accession No. CRL-11537), for the human Y4 receptor, L-hY4-3 (deposited January 11, 1995, under ATCC Accession No. CRL-11779), and for the human Y5 receptor, L-hY5-7 (deposited November 15, 1995, under ATCC Accession No. CRL-11995).

Expression of other G-protein coupled receptors

**$\alpha_1$  Human Adrenergic Receptors:** To determine the binding of compounds to human  $\alpha_1$  receptors, LM(tk-) cell lines stably transfected with the genes encoding the  $\alpha_{1a}$ ,  $\alpha_{1b}$ , and  $\alpha_{1d}$  receptors were used. The nomenclature describing the  $\alpha_1$  receptors was changed recently, such that the receptor formerly designated  $\alpha_{1a}$  is now designated  $\alpha_{1d}$ , and the receptor formerly designated  $\alpha_{1c}$  is now designated  $\alpha_{1a}$  (ref). The cell lines expressing these receptors were deposited with the ATCC before the nomenclature change and reflect the subtype designations formerly assigned to these receptors. Thus, the cell line expressing the

receptor described herein as the  $\alpha_{1a}$  receptor was deposited with the ATCC on September 25, 1992, under ATCC Accession No. CRL 11140 with the designation L- $\alpha_{1c}$ . The cell line expressing receptor described herein as the  $\alpha_{1a}$  receptor was deposited with the ATCC on September 25, 1992, under ATCC Accession No. CRL 11138 with the designation L- $\alpha_{1A}$ . The cell line expressing the  $\alpha_{1b}$  receptor is designated L- $\alpha_{1B}$ , and was deposited on September 25, 1992, under ATCC Accession No. CRL 11139.

**$\alpha_2$  Human Adrenergic Receptors:** To determine the binding of compounds to human  $\alpha_2$  receptors, LM(tk-) cell lines stably transfected with the genes encoding the  $\alpha_{2A}$ ,  $\alpha_{2B}$ , and  $\alpha_{2C}$  receptors were used. The cell line expressing the  $\alpha_{2A}$  receptor is designated L- $\alpha_{2A}$ , and was deposited on November 6, 1992, under ATCC Accession No. CRL 11180. The cell line expressing the  $\alpha_{2B}$  receptor is designated L-NGC- $\alpha_{2B}$ , and was deposited on October 25, 1989, under ATCC Accession No. CRL 10275. The cell line expressing the  $\alpha_{2C}$  receptor is designated L- $\alpha_{2C}$ , and was deposited on November 6, 1992, under ATCC Accession No. CRL-11181. Cell lysates were prepared as described below (see Radioligand Binding to Membrane Suspensions), and suspended in 25mM glycylglycine buffer (pH 7.6 at room temperature). Equilibrium competition binding assay were performed using [ $^3$ H]rauwolscine (0.5nM), and nonspecific binding was determined by incubation with 10 $\mu$ M phentolamine. The bound radioligand was separated by filtration through GF/B filters using a cell harvester.

**Human Histamine  $H_1$  Receptor:** The coding sequence of the human histamine  $H_1$  receptor, homologous to the bovine  $H_1$  receptor, was obtained from a human hippocampal cDNA library, and was cloned into the eukaryotic expression vector pCEXV-3. The plasmid DNA for the  $H_1$  receptor is

-93-

designated pcEXV-H1, and was deposited on November 6, 1992, under ATCC Accession No. 75346. This construct was transfected into COS-7 cells by the DEAE-dextran method. Cells were harvested after 72 hours and lysed by sonication in 5mM Tris-HCl, 5mM EDTA, pH 7.5. The cell lysates were centrifuged at 1000 rpm for 5 min at 4°C, and the supernatant was centrifuged at 30,000 x g for 20 min. at 4°C. The pellet was suspended in 37.8 mM NaHPO<sub>4</sub>, 12.2 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.5. The binding of the histamine H<sub>1</sub> antagonist [<sup>3</sup>H]mepyramine (1nM, specific activity: 24.8 Ci/mM) was done in a final volume of 0.25 mL and incubated at room temperature for 60 min. Nonspecific binding was determined in the presence of 10 μM mepyramine. The bound radioligand was separated by filtration through GF/B filters using a cell harvester.

**Human Histamine H<sub>2</sub> Receptor:** The coding sequence of the human H<sub>2</sub> receptor was obtained from a human placenta genomic library, and cloned into the cloning site of PCEXV-3 eukaryotic expression vector. The plasmid DNA for the H<sub>2</sub> receptor is designated pcEXV-H2, and was deposited on November 6, 1992 under ATCC Accession No. 75345. This construct was transfected into COS-7 cells by the DEAE-dextran method. Cells were harvested after 72 hours and lysed by sonication in 5mM Tris-HCl, 5mM EDTA, pH 7.5. The cell lysates were centrifuged at 1000 rpm for 5 min at 4°C, and the supernatant was centrifuged at 30,000 x g for 20 min at 4 °C. The pellet was suspended in 37.8 mM NaHPO<sub>4</sub>, 12.2 mM K<sub>2</sub>PO<sub>4</sub>, pH 7.5. The binding of the histamine H<sub>2</sub> antagonist [<sup>3</sup>H]tiotidine (5nM, specific activity: 70 Ci/mM) was done in a final volume of 0.25 ml and incubated at room temperature for 60 min. Nonspecific binding was determined in the presence of 10 μM histamine. The bound radioligand was separated by filtration through GF/B filters using a cell harvester.

**Human Serotonin Receptors:**

5HT<sub>1D $\alpha$</sub> , 5HT<sub>1D $\beta$</sub> , 5HT<sub>1E</sub>, 5HT<sub>1F</sub> Receptors: LM(tk-) clonal cell lines stably transfected with the genes encoding each of these 5HT receptor subtypes were prepared as described above. The cell line for the 5HT<sub>1D $\alpha$</sub>  receptor, designated as Ltk-8-30-84, was deposited on April 17, 1990, and accorded ATCC Accession No. CRL 10421. The cell for the 5HT<sub>1D $\beta$</sub>  receptor, designated as Ltk-11, was deposited on April 17, 1990, and accorded ATCC Accession No. CRL 10422. The cell line for the 5HT<sub>1E</sub> receptor, designated 5 HT<sub>1E</sub>-7, was deposited on November 6, 1991, and accorded ATCC Accession No. CRL 10913. The cell line for the 5HT<sub>1F</sub> receptor, designated L-5-HT<sub>1F</sub>, was deposited on December 27, 1991, and accorded ATCC Accession No. ATCC 10957. Membrane preparations comprising these receptors were prepared as described below, and suspended in 50mM Tris-HCl buffer (pH 7.4 at 37°C) containing 10 mM MgCl<sub>2</sub>, 0.2 mM EDTA, 10 $\mu$ M pargyline, and 0.1% ascorbate. The binding of compounds was determined in competition binding assays by incubation for 30 minutes at 37°C in the presence of 5nM [<sup>3</sup>H]serotonin. Nonspecific binding was determined in the presence of 10 $\mu$ M serotonin. The bound radioligand was separated by filtration through GF/B filters using a cell harvester.

**Human 5HT<sub>2</sub> Receptor:** The coding sequence of the human 5HT<sub>2</sub> receptor was obtained from a human brain cortex cDNA library, and cloned into the cloning site of pCEXV-3 eukaryotic expression vector. This construct was transfected into COS-7 cells by the DEAE-dextran method. Cells were harvested after 72 hours and lysed by sonication in 5mM Tris-HCl, 5mM EDTA, pH 7.5. This cell line was deposited with the ATCC on October 31, 1989, designated as L-NGC-5HT<sub>2</sub>, and was accorded ATCC Accession No. CRL 10287. The cell lysates were centrifuged at 1000



-95-

rpm for 5 minutes at 4°C, and the supernatant was centrifuged at 30,000 x g for 20 minutes at 4°C. The pellet was suspended in 50mM Tris-HCl buffer (pH 7.7 at room temperature) containing 10 mM MgSO<sub>4</sub>, 0.5mM EDTA, and 0.1% ascorbate. The potency of alpha-1 antagonists at 5HT<sub>2</sub> receptors was determined in equilibrium competition binding assays using [3H]ketanserin (1nM). Nonspecific binding was defined by the addition of 10μM mianserin. The bound radioligand was separated by filtration through GF/B filters using a cell harvester.

**Human 5-HT<sub>2</sub> Receptor:** A LM(tk-) clonal cell line stably transfected with the gene encoding the 5HT<sub>2</sub> receptor subtype was prepared as described above. The cell line for the 5HT<sub>2</sub> receptor, designated as L-5HT<sub>2B</sub>, was deposited on October 20, 1992, and accorded ATCC Accession No. CRL 11166.

**Human Dopamine D<sub>3</sub> Receptor:** The binding of compounds to the human D<sub>3</sub> receptor was determined using membrane preparations from COS-7 cells transfected with the gene encoding the human D<sub>3</sub> receptor. The human dopamine D<sub>3</sub> receptor was prepared according to known methods (Sokoloff, P. et al. Nature, 347, 146, 1990, deposited with the EMBL Genbank as X53944). Cells were harvested after 72 hours and lysed by sonication in 5mM Tris-HCl, 5mM EDTA, pH 7.5. The cell lysates were centrifuged at 1000 rpm for 5 minutes at 4°C, and the supernatant was centrifuged at 30,000 x g for 20 minutes at 4°C. The pellet was suspended in 50 mM Tris-HCl (pH 7.4) containing 1mM EDTA, 5mM KCl, 1.5mM CaCl<sub>2</sub>, 4mM MgCl<sub>2</sub>, and 0.1% ascorbic acid. The cell lysates were incubated with [<sup>3</sup>H]spiperone (2nM), using 10μM (+)Butaclamol to determine nonspecific binding.

Membrane Harvest

Membranes were harvested from COS-7 cells 48 hours after transient transfection. Adherent cells were washed twice in ice-cold phosphate buffered saline (138 mM NaCl, 8.1 mM Na<sub>2</sub>HPO<sub>4</sub>, 2.5 mM KCl, 1.2 mM KH<sub>2</sub>PO<sub>4</sub>, 0.9 mM CaCl<sub>2</sub>, 0.5 mM MgCl<sub>2</sub>, pH 7.4) and lysed by sonication in ice-cold sonication buffer (20 mM Tris-HCl, 5 mM EDTA, pH 7.7). Large particles and debris were cleared by low speed centrifugation (200 x g, 5 min, 4 °C). Membranes were collected from the supernatant fraction by centrifugation (32,000 x g, 18 min, 4 °C), washed with ice-cold hypotonic buffer, and collected again by centrifugation (32,000 x g, 18 min, 4 °C). The final membrane pellet was resuspended by sonication into a small volume of ice-cold binding buffer (~1 ml for every 5 plates: 10 mM NaCl, 20 mM HEPES, 0.22 mM KH<sub>2</sub>PO<sub>4</sub>, 1.26 mM CaCl<sub>2</sub>, 0.81 mM MgSO<sub>4</sub>, pH 7.4). Protein concentration was measured by the Bradford method (Bradford, 1976) using Bio-Rad Reagent, with bovine serum albumin as a standard. Membranes were held on ice for up to one hour and used fresh, or flash-frozen and stored in liquid nitrogen.

Membranes were prepared similarly from 293, LM(tk-), and NIH-3T3 cells. To prepare membranes from baculovirus infected cells, 2 x 10<sup>7</sup> Sf21 cells were grown in 150mm tissue culture dishes and infected with a high-titer stock of hY5BB3. Cells were incubated for 2-4 days at 27°C, no CO<sub>2</sub> before harvesting and membrane preparation as described above.

Membranes were prepared similarly from dissected rat hypothalamus. Frozen hypothalami were homogenized for 20 seconds in ice-cold sonication buffer with the narrow probe of a Virtishear homogenizer at 1000 rpm (Virtis, Gardiner, NY). Large particles and debris were cleared by

-97-

centrifugation (200 x g, 5 min, 4 °C) and the supernatant fraction was reserved on ice. Membranes were further extracted from the pellet by repeating the homogenization and centrifugation procedure two more times. The supernatant fractions were pooled and subjected to high speed centrifugation (100,000 x g, 20 min. 4 °C). The final membrane pellet was resuspended by gentle homogenization into a small volume of ice-cold binding buffer (1 mL/ gram wet weight tissue) and held on ice for up to one hour, or flash-frozen and stored in liquid nitrogen.

#### Radioligand Binding to Membrane Suspensions

Membrane suspensions were diluted in binding buffer supplemented with 0.1% bovine serum albumin to yield an optimal membrane protein concentration so that  $^{125}\text{I}$ -PYY (or alternative radioligand such as  $^{125}\text{I}$ -NPY,  $^{125}\text{I}$ -PYY<sub>3-36</sub>, or  $^{125}\text{I}$ -[Leu<sup>31</sup>Pro<sup>34</sup>]PYY) bound by membranes in the assay was less than 10% of  $^{125}\text{I}$ -PYY (or alternative radioligand) delivered to the sample (100,000 dpm/sample = 0.08 nM for competition binding assays).  $^{125}\text{I}$ -PYY (or alternative radioligand) and peptide competitors were also diluted to desired concentrations in supplemented binding buffer. Individual samples were then prepared in 96-well polypropylene microtiter plates by mixing  $^{125}\text{I}$ -PYY (25  $\mu\text{L}$ ) (or alternative radioligand), competing peptides or supplemented binding buffer (25  $\mu\text{L}$ ), and finally, membrane suspensions (200  $\mu\text{L}$ ). Samples were incubated in a 30 °C water bath with constant shaking for 120 min. Incubations were terminated by filtration over Whatman GF/C filters (pre-coated with 1% polyethyleneimine and air-dried before use), followed by washing with 5 mL of ice-cold binding buffer. Filter-trapped membranes were impregnated with MultiLex solid scintillant (Wallac, Turku, Finland) and counted for  $^{125}\text{I}$  in a Wallac Beta-

Plate Reader. Non-specific binding was defined by 300 nM human NPY for all receptors except the Y4 subtypes; 100 nM human PP was used for the human Y4 and 100 nM rat PP for the rat Y4. Specific binding in time course and competition studies was typically 80%; most non-specific binding was associated with the filter. Binding data were analyzed using nonlinear regression and statistical techniques available in the GraphPAD Prism package (San Diego, CA).

#### Functional Assay: Radioimmunoassay of cAMP

Stably transfected cells were seeded into 96-well microtiter plates and cultured until confluent. To reduce the potential for receptor desensitization, the serum component of the media was reduced to 1.5% for 4 to 16 hours before the assay. Cells were washed in Hank's buffered saline, or HBS (150 mM NaCl, 20 mM HEPES, 1 mM CaCl<sub>2</sub>, 5 mM KCl, 1 mM MgCl<sub>2</sub>, and 10 mM glucose) supplemented with 0.1% bovine serum albumin plus 5 mM theophylline and pre-equilibrated in the same solution for 20 min at 37 °C in 5% CO<sub>2</sub>. Cells were then incubated 5 min with 10 μM forskolin and various concentrations of receptor-selective ligands. The assay was terminated by the removal of HBS and acidification of the cells with 100 mM HCl. Intracellular cAMP was extracted and quantified with a modified version of a magnetic bead-based radioimmunoassay (Advanced Magnetix, Cambridge, MA). The final antigen/antibody complex was separated from free <sup>125</sup>I-cAMP by vacuum filtration through a PVDF filter in a microtiter plate (Millipore, Bedford, MA). Filters were punched and counted for <sup>125</sup>I in a Packard gamma counter. Binding data were analyzed using nonlinear regression and statistical techniques available in the GraphPAD Prism package (San Diego, CA).

Functional Assay: Intracellular calcium mobilization

The intracellular free calcium concentration was measured by microspectrofluorimetry using the fluorescent indicator dye Fura-2/AM (ref). Stably transfected cells were seeded onto a 35 mm culture dish containing a glass coverslip insert. Cells were washed with HBS and loaded with 100  $\mu$ l of Fura-2/AM (10  $\mu$ M) for 20 to 40 min. After washing with HBS to remove the Fura-2/AM solution, cells were equilibrated in HBS for 10 to 20 min. Cells were then visualized under the 40X objective of a Leitz Fluovert FS microscope and fluorescence emission was determined at 510 nm with excitation wave lengths alternating between 340 nm and 380 nm. Raw fluorescence data were converted to calcium concentrations using standard calcium concentration curves and software analysis techniques.

In vivo STUDIES IN RATS

20

Food intake in satiated rats

For these determinations food intake maybe measured in normal satiated rats after intracerebroventricular application (i.c.v.) of NPY in the presence or absence of the test compound. Male Sprague Dawley rats ciba-Geigy AG, Sisseln, Switzerland weighing between 180g and 220 g are used for all experiments. The rats are individually housed in stainless steel cages and maintained on an 11:13 h light-dark cycle (lights off at 18:00 h) at a controlled temperature of 21-23 °C at all times. Water and food (NAFAG lab chow pellets, NAFAG, Gossau, Switzerland) are available ad libidum.

Rats under pentobarbital anesthesia are stereotaxically implanted with a stainless steel guide cannula targeted

-100-

at the right lateral ventricle. Stereotaxic coordinates, with the incisor bar set -2.0mm below interaural line, are: -0.8mm anterior and +1.3mm lateral to bregma. The guide cannula is placed on the dura. Injection cannulas extend the guide cannulas -3.8mm ventrally to the skull surface. Animals are allowed at least 4 days of recovery postoperatively before being used in the experiments. Cannula placement is checked postoperatively by testing all rats for their drinking response to a 50 ng intracerebroventricular (i.c.v.) injection of angiotensin II. Only rats which drink at least 2.5 ml of water within 30 min. after angiotensin II injection are used in the feeding studies.

All injections are made in the morning 2 hours after light onset. Peptides are injected in artificial cerebrospinal fluid (ACSF) in a volume of 5 $\mu$ l. ACSF contains: NaCl 124mM, KCl 3.75 mM, CaCl<sub>2</sub> 2.5 mM, MgSO<sub>4</sub> 2.0 mM, KH<sub>2</sub>PO<sub>4</sub> 0.22mM, NaHCO<sub>3</sub> 26 mM and glucose 10 mM. porcine-NPY is dissolved in artificial cerebrospinal fluid (ACS). For i.c.v. injection the test compounds are preferably dissolved in DMSO/water (10%, v/v). The vehicle used for intraperitoneal (i.p.), subcutaneous (s.c.) or oral (p.o.) delivery of compounds is preferably water, physiological saline or DMSO/water (10% v/v), or cremophor/water (20% v/v), respectively.

Animals which are treated with both test compounds and p-NPY are treated first with the test compound. Then, 10 min. after i.c.v. application of the test compound or vehicle (control), or 30-60 min after i.p., s.c. and p.o. application of the test compound or vehicle, 300 pmol of NPY is administered by intracerebroventricular (i.c.v.) application.

35

-101-

Food intake may be measured by placing preweighed pellets into the cages at the time of NPY injection. Pellets are removed from the cage subsequently at each selected time point and replaced with a new set of preweighed pellets.

5 The food intake of animals treated with test compound may be calculated as a percentage of the food intake of control animals, i.e., animals treated with vehicle. Alternatively, food intake for a group of animals subjected to the same experimental condition may be

10 expressed as the mean  $\pm$  S.E.M. Statistical analysis is performed by analysis of variance using the Student-Newman-Keuls test.

#### Food intake in food-deprived rats

15 Food-deprivation experiments are conducted with male Sprague-Dawley rats weighing between 220 and 250 g. After receipt, the animals are individually housed for the duration of the study and allowed free access to normal food together with tap water. The animals are

20 maintained in a room with a 12 h light/dark cycle (8:00 a.m. to 8:00 p.m. light) at 24 °C and monitored humidity. After placement into individual cages the rats undergo a 4 day equilibration period, during which they are habituated to their new environment and to eating a

25 powdered or pellet diet (NAFAG, Gossau, Switzerland).

At the end of the equilibration period, food is removed from the animals for 24 hours starting at 8:00 a.m. At the end of the fasting period compound or vehicle may be

30 administered to the animals orally or by injection intraperitoneally or intravenously. After 10 - 60 min. food is returned to the animals and their food intake monitored at various time periods during the following 24 hour period. The food intake of animals treated with

35 test compound may be calculated as a percentage of the

food intake of control animals (i.e., animals treated with vehicle). Alternatively, food intake for a group of animals subjected to the same experimental conditions may be expressed as the mean  $\pm$  S.E.M.

5

#### Food intake in obese Zucker rats

The antiobesity efficacy of the compounds according to the present invention might also be manifested in Zucker obese rats, which are known in the art as an animal model of obesity. These studies are conducted with male Zucker fatty rats (fa/fa Harlan CPB, Austerlitz NL) weighing between 480g and 500g. Animals are individually housed in metabolism cages for the duration of the study and allowed free access to normal powdered food and water. The animals are maintained in a room with a 12 h light/dark cycle (light from 8:00 A.M. to 8:00 P.M.) at 24°C and monitored humidity. After placement into the metabolism cages the rats undergo a 6 day equilibration period, during which they are habituated to their new environment and to eating a powdered diet. At the end of the equilibration period, food intake during the light and dark phases is determined. After a 3 day control period, the animals are treated with test compounds or vehicle (preferably water or physiological saline or DMSO/water (10%,v/v) or cremophor/water (20%,v/v)). Food intake is then monitored over the following 3 day period to determine the effect of administration of test compound or vehicle alone. As in the studies described hereinabove, food intake in the presence of drug may be expressed as a percentage of the food intake of animals treated with vehicle.

#### Materials

Cell culture media and supplements were from Specialty Media (Lavallette, NJ). Cell culture plates (150 mm and

35



-103-

96-well microtiter) were from Corning (Corning, NY). Sf9, Sf21, and High Five insect cells, as well as the baculovirus transfer plasmid, pBlueBacIII™, were purchased from Invitrogen (San Diego, CA). TMN-FH insect  
5 medium complemented with 10% fetal calf serum, and the baculovirus DNA, BaculoGold™, was obtained from Pharmingen (San Diego, CA.). Ex-Cell 400™ medium with L-Glutamine was purchased from JRH Scientific. Polypropylene 96-well microtiter plates were from Co-star  
10 (Cambridge, MA). All radioligands were from New England Nuclear (Boston, MA). Commercially available NPY and related peptide analogs were either from Bachem California (Torrance, CA) or Peninsula (Belmont, CA); [D-Trp<sup>32</sup>]NPY and PP C-terminal fragments were synthesized by  
15 custom order from Chiron Mimotopes Peptide Systems (San Diego, CA). Bio-Rad Reagent was from Bio-Rad (Hercules, CA). Bovine serum albumin (ultra-fat free, A-7511) was from Sigma (St. Louis, MO). All other materials were reagent grade.

## EXPERIMENTAL RESULTS

Applicants have synthesized and evaluated the binding and functional properties of several compounds at the cloned human Y1, human Y2, human Y4, and human Y5 receptors. As shown below in Table 5, applicants have discovered several compounds which not only bind selectively to the human Y5 receptor but also act as Y5 receptor antagonists, as measured by their ability to block NPY-induced inhibition of cAMP accumulation in forskolin-stimulated LM(tk-) cells stably transfected with the cloned human Y5 receptor.

**Table 5: Evaluation of human Y5 receptor antagonists**

The ability of the compounds to antagonize the Y-type receptors is reported as the  $K_b$ . The  $K_b$  is derived from the  $EC_{50}$ , or concentration of half-maximal effect, in the presence ( $EC_{50}$ ) or absence ( $EC_{50}'$ ) of compound, according to the equation:  $K_b = [NPY] / ((EC_{50}/EC_{50}') - 1)$ . Results shown are representative of at least three independent experiments.

N.D. = Not determined.

**Table 5**

|         | Binding Affinity<br>(K <sub>i</sub> (nM) vs. <sup>125</sup> I-PYY) |      |       |    |                     |
|---------|--|------|-------|----|---------------------|
| Example | Human Receptor   |      |       |    | K <sub>b</sub> (nM) |
| -       | Y1   | Y2   | Y4    | Y5 | -                   |
| 31      | 5550   | 1000 | 8020  | 14 | 6.0                 |
| 32      | 3550   | 955  | 11700 | 11 | 23                  |

-105-

|    | Binding Affinity<br>(K <sub>i</sub> (nM) vs. <sup>125</sup> I-PYY) |       |       |     |     |
|----|--|-------|-------|-----|-----|
| 36 | 16000  | 7760  | 20400 | 8.3 | 26  |
| 38 | 13000  | 1610  | 18500 | 9.8 | 16  |
| 40 | 17200  | 7570  | 27500 | 11  | 3.0 |
| 37 | 14500  | 617   | 21500 | 26  | 38  |
| 77 | 3240   | 851   | 13100 | 17  | 311 |
| 44 | 23700  | 58200 | 19300 | 14  | 50  |
| 45 | 48700  | 5280  | 63100 | 28  | 49  |

10 Several of the compounds were further tested using in  
15 vivo animal models of feeding behavior. Since NPY is the  
strongest known stimulant of feeding behavior,  
experiments were performed with several compounds to  
evaluate the effect of the compounds described above on  
NPY-induced feeding behavior in satiated rats.

First, 300 pmole of porcine NPY in vehicle (A.C.S.F.) was  
administered by intracerebroventricular (i.c.v.)  
injection, along with i.p. administration of compound  
20 vehicle (10% DMSO/water), and the food intake of NPY-  
stimulated animals was compared to food intake in animals  
treated with the vehicles. The 300 pmole injection of  
NPY was found to significantly induce food intake ( $p <$   
0.05; Student-Newman-Keuls).

25 Using the 300 pmole dose of NPY found to be effective to  
stimulate feeding, other animals were treated with the  
compounds by intraperitoneal (i.p.) administration,

-106-

followed 30-60 minutes later by i.c.v. NPY administration, and measurement of subsequent food intake. As shown in Table 6, NPY-induced food intake was significantly reduced in animals first treated with the compounds ( $p < 0.05$ ; Student-Newman-Keuls). These experiments demonstrate that NPY-induced food intake is significantly reduced by administration to animals of a compound which is a Y5-selective antagonist.

Table 6. NPY-induced cumulative food intake in rats treated with either the i.c.v. and i.p. vehicles (control), 300 pmole NPY alone (NPY), or in rats treated first with compound and then NPY (NPY + compound). Food intake was measured 4 hours after stimulation with NPY. Food intake is reported as the mean  $\pm$  S.E.M. intake for a group of animals.

Table 6

| Example                       | 31            | 32            |
|-------------------------------|---------------|---------------|
| Compound Dose<br>(mg/kg i.p.) | 10            | 30            |
|                               |               |               |
| control<br>(vehicles<br>only) | 2.4 $\pm$ 0.7 | 2.9 $\pm$ 0.8 |
| NPY                           | 5.8 $\pm$ 0.5 | 4.9 $\pm$ 0.4 |
| NPY +<br>compound             | 3.8 $\pm$ 0.4 | 1.5 $\pm$ 0.6 |

-107-

Since food deprivation induces an increase in the hypothalamic NPY levels, it has been postulated that food intake following a period of food deprivation is NPY-mediated. Therefore, the Y5 antagonists of Table 5 were administered to conscious rats following a 24h food deprivation. Each of the human Y5 receptor antagonists shown in Table 5 was able to significantly reduce NPY-induced food intake in the animals, as shown below in Table 7. The food intake of animals treated with test compound is reported as a percentage of the food intake measured for control animals (treated with vehicle), i.e., 25% means the animals treated with the compound consumed only 25% as much food as the control animals. Measurements were performed two hours after administration of the test compound.

Table 7 Two-hour food intake of NPY-stimulated rats. Food intake is expressed as the percentage of intake compared to control rats.

|    |  |         |          |
|----|--|---------|----------|
| 5  |  | Example | Mean (%) |
|    |  | 31      | 27       |
|    |  | 32      | 36       |
|    |  | 36      | 35       |
|    |  | 38      | 80       |
| 10 |  | 40      | 55       |
|    |  | 37      | 58       |
|    |  | 77      | 32       |
|    |  | 44      | 73       |
|    |  | 45      | 84       |

15

These experiments indicate that the compounds of the present invention inhibit food intake in rats, especially when administered in a range of about 0.01 to about 100 mg/kg rat, by either oral, intraperitoneal or intravenous administration. The animals appeared normal during these experiments, and no ill effects on the animals were observed after the termination of the feeding experiments.

20

25

The binding properties of the compounds were also evaluated with respect to other cloned human G-protein coupled receptors. As shown in Table 8, below, the Y5-

selective compounds described hereinabove exhibited lower affinity for receptors other than the Y-type receptors.

Table 8 Cross-reactivity of compounds at other cloned human receptors

[illegible]

Table 8 (continued)

| Compound | Receptor (pKi)    |                  |                  |                   |                   |                    |                    |
|----------|-------------------|------------------|------------------|-------------------|-------------------|--------------------|--------------------|
|          | 5HT <sub>1a</sub> | 5HT <sub>2</sub> | 5HT <sub>7</sub> | 5HT <sub>1F</sub> | 5HT <sub>1E</sub> | 5HT <sub>1Dβ</sub> | 5HT <sub>1Dα</sub> |
| 31       | 5.88              | 6.74             | 6.50             | 5.30              | 5.30              | 5.30               | 5.32               |
| 32       | 5.54              | 6.55             | 6.42             | 5.30              | 5.30              | 5.30               | 6.04               |
| 36       | 6.73              | 5.93             | 6.37             | 5.30              | 5.30              | 5.37               | 5.94               |
| 38       | N.D.              | N.D.             | N.D.             | N.D.              | N.D.              | N.D.               | N.D.               |
| 40       | 6.56              | 5.99             | 6.39             | 5.30              | 5.30              | 5.41               | 5.98               |
| 37       | N.D.              | N.D.             | N.D.             | N.D.              | N.D.              | N.D.               | N.D.               |
| 77       | 5.82              | 5.99             | 5.35             | 5.30              | 5.30              | 5.39               | 5.62               |
| 44       | N.D.              | N.D.             | N.D.             | N.D.              | N.D.              | N.D.               | N.D.               |
| 45       | N.D.              | N.D.             | N.D.             | N.D.              | N.D.              | N.D.               | N.D.               |

5

10



-111-

EXPERIMENTAL DISCUSSION

Y5 receptors are highly attractive targets for appetite and weight control based on several lines of research (Sahu and Kalra, 1993). NPY is the most potent stimulant of feeding behavior yet described (Clark et al., 1984; Levine and Morley, 1984; Stanley and Leibowitz, 1984). Direct injection of NPY into the hypothalamus of rats can increase food intake ~ 10-fold over a 4-hour period (Stanley et al., 1992). NPY-stimulated rats display a preference for carbohydrates over protein and fat (Stanley et al., 1985). Interestingly, NPY and NPY mRNA are increased in food-deprived rats (Brady et al., 1990; O' Shea and Gundlach, 1991) and also in rats which are genetically obese (Sanacora et al., 1990) or made diabetic by treatment with streptozotocin (White et al., 1990). One potential explanation is that NPY, a potent stimulant of feeding behavior in normal rats, is disregulated in the overweight or diabetic animal so that food intake is increased, accompanied by obesity. The physiological stress of obesity increases the risk for health problems such as cardiovascular malfunction, osteoarthritis, and hyperinsulinemia, together with a worsened prognosis for adult-onset diabetes. A nonpeptide antagonist targeted to the Y5 receptor could therefore be effective as a way to control not only appetite and body weight but an entire range of obesity- and diabetes-related disorders (Dryden et al., 1994). There is also neurochemical evidence to suggest that NPY-mediated functions are disregulated in eating disorders such as bulimia and anorexia nervosa, so that they too could be responsive to treatment by a Y5-selective drug. It has been proposed, for example, that food intake in NPY-stimulated rats mimics the massive food consumption associated with binge eating in bulimia (Stanley, 1993).

-112-

CSF levels of PYY but not NPY were elevated in bulimic patients who abstained from bingeing, and then diminished when bingeing was allowed (Berrettini et al., 1988). Conversely, NPY levels were elevated in underweight anorectic patients and then diminished as body weight was normalized (Kaye et al., 1990).

As described above, the human and rat *in vitro* expression models were used in combination to screen for compounds intended to modulate NPY-dependent feeding behavior. Using this approach, applicants have discovered several compounds which inhibit feeding behavior in animal models, which should lead to additional drug discoveries. The compounds according to the present invention inhibit food intake in Zucker obese rats in a range especially of about 0.01 to about 100 mg/kg after oral, intraperitoneal or intravenous administration.

The Y5 pharmacological profile further offers a new standard by which to review the molecular basis of all NPY-dependent processes. Such an exercise suggests that the Y5 receptor is likely to have a physiological significance beyond feeding behavior. It has been reported, for example, that a Y-type receptor can regulate luteinizing hormone releasing hormone (LHRH) release from the median eminence of steroid-primed rats *in vitro* with an atypical Y1 pharmacological profile. NPY, NPY<sub>7-16</sub>, and LP-NPY were all effective at 1 $\mu$ M but deletion of as few as four amino acids from the N-terminus of NPY destroyed biological activity. The Y5 may therefore represent a therapeutic target for sexual or reproductive disorders. It is worth while considering that the Y5 is so similar in pharmacological profile to the other Y-type receptors that it may have been

-113-

overlooked among a mixed population of Y1, Y2 and Y4 receptors. Certain functions now associated with these subtypes could therefore be reassigned to Y5 as our pharmacological tools grow more sophisticated. By offering new insight into NPY receptor pharmacology, the Y5 thereby provides a greater clarity and focus in the field of drug design.

TABLE 9: Pathophysiological Conditions Associated With NPY

|    |  |   |                              |
|----|--|---|------------------------------|
| 5  | The following pathological conditions have been linked to either 1) application of exogenous NPY, or 2) changes in levels of endogenous NPY. |   |                              |
|    | 1  | obesity   | Sahu and Kalra, 1993         |
|    | 2  | eating disorders (anorexia and bulimia nervosa) | Stanley, 1993                |
| 10 | 3  | sexual/reproductive function                    | Clark, 1994                  |
|    | 4  | depression                                      | Heilig and Weiderlov, 1990   |
|    | 5  | anxiety   | Wahlestedt et al., 1993      |
|    | 6  | cocaine addiction                               | Wahlestedt et al., 1991      |
|    | 7  | gastric ulcer                                   | Penner et al., 1993          |
| 15 | 8  | memory loss                                     | Morley and Flood, 1990       |
|    | 9  | pain  | Hua et al., 1991             |
|    | 10   | epileptic seizure                               | Rizzi et al., 1993           |
|    | 11   | hypertension                                    | Zukowska-Grojec et al., 1993 |
|    | 12   | subarachnoid hemorrhage                         | Abel et al., 1988            |
| 20 | 13   | shock   | Hauser et al., 1993          |
|    | 14   | circadian rhythm                                | Albers and Ferris, 1984      |
|    | 15   | nasal congestion                                | Lacroix et al., 1988         |
|    | 16   | diarrhea  | Cox and Cuthbert, 1990       |
| 25 | 17   | neurogenic voiding dysfunction                  | Zoubek et al., 1993          |

-115-

A successful strategy for the design of a Y5-receptor based drug or for any drug targeted to single G protein-coupled receptor subtype involves the screening of candidate compounds 1) in radioligand binding assays so as to detect affinity for cross-reactive G protein-coupled receptors, and 2) in physiological assays so as to detect undesirable side effects. In the specific process of screening for a Y5-selective drug, the receptor subtypes most likely to cross-react and therefore most important for radioligand binding screens include the other "Y-type" receptors, Y1, Y2, Y3, and Y4. Cross-reactivity between the Y5 and any of the other subtypes could result in potential complications as suggested by the pathophysiological indications listed in Table 9. In designing a Y5 antagonist for obesity and appetite control, for example, it is important not to design a Y1 antagonist resulting in hypertension or increased anxiety, a Y2 antagonist resulting in memory loss, or a Y4 antagonist resulting in increased appetite.

TABLE 10: Y-Type Receptor Indications

|    | Y-type<br>Receptor<br>Indications                         | Receptor<br>Subtype                       | Drug<br>Activity                    | Reference   |
|----|---|---|-------------------------------------|---|
| 5  | obesity,<br>appetite<br>disorder                          | atypical Y1                               | antagonist                          | Sahu and<br>Kalra,<br>1993  |
| 10 | adult onset<br>diabetes                                   | atypical Y1                               | antagonist                          | Sahu and<br>Kalra,<br>1993  |
|    | bulimia<br>nervosa  | atypical Y1                               | antagonist                          | Stanley,<br>1993  |
| 15 | pheochromoc<br>ytoma-<br>induced<br>hypertensio<br>n      | Y1  | antagonist                          | Grouzman<br>et al.,<br>1989   |
| 20 | subarachnoi<br>d<br>hemorrhage                            | Y1  | antagonist                          | Abel et<br>al., 1988  |
|    | neurogenic<br>vascular<br>hypertrophy                     | Y1<br>Y2                                  | antagonist<br>antagonist            | Zukowska-<br>Grojec et<br>al., 1993                                 |
| 25 | epileptic<br>seizure                                      | Y2  | antagonist                          | Rizzi et<br>al., 1993   |
| 30 | hypertensio<br>n:<br>central,<br>peripheral<br>regulation | peripheral Y1<br>central Y3<br>central Y2 | antagonist<br>agonist<br>antagonist | Grundemar<br>and<br>Hakanson,<br>1993<br>Barraco<br>et al.,<br>1991 |
|    | obesity,<br>appetite<br>disorder                          | Y4 or PP                                  | agonist                             | Malaisse-<br>Lagae et<br>al., 1977                                  |
| 35 | anorexia<br>nervosa                                       | atypical Y1                               | agonist                             | Berrettin<br>i et al.,<br>1988                                      |

-117-

|    |                              |                |                    |                         |
|----|------------------------------|----------------|--------------------|-------------------------|
|    | anxiety                      | Y1             | agonist            | Wahlestedt et al., 1993 |
|    | cocaine addiction            | Y1             | agonist            | Wahlestedt et al., 1991 |
| 5  | stress-induced gastric ulcer | Y1<br>Y4 or PP | agonist<br>agonist | Penner et al., 1993     |
|    | memory loss                  | Y2             | agonist            | Morley and Flood, 1990  |
|    | pain                         | Y2             | agonist            | Hua et al., 1991        |
| 10 | shock                        | Y1             | agonist            | Hauser et al., 1993     |
|    | sleep disturbances, jet lag  | Y2             | not clear          | Albers and Ferris, 1984 |
| 15 | nasal decongestion           | Y1<br>Y2       | agonist<br>agonist | Lacroix et al., 1988    |
|    | diarrhea                     | Y2             | agonist            | Cox and Cuthbert, 1990  |

-118-

The Y5 receptor represents an enormous opportunity for the development of novel and selective drug therapies, particularly those targeted to appetite and weight control, but also for memory loss, depression, anxiety, gastric ulcer, epileptic seizure, pain, hypertension, subarachnoid hemorrhage, sleeping disturbances, nasal congestion, neurogenic voiding dysfunction, and diarrhea.

In particular, the discovery of Y5-selective antagonists which inhibit food intake in rats provides a method of modifying feeding behavior in a wide variety of vertebrate animals.



REFERENCES

- Abel, P.W., Han, C., Noe, B.D., and McDonald, J.K.  
(1988). Neuropeptide Y: vasoconstrictor effects and  
5 possible role in cerebral vasospasm after experimental  
subarachnoid hemorrhage. Brain Res. 463: 250-258.
- Albers, H.E., and Ferris, C.F. (1984). Neuropeptide Y:  
Role in light-dark cycle entrainment of hamster circadian  
10 rhythms. Neurosci. Lett. 50: 163-168.
- Aruffo, A. and Seed, B. (1987). Molecular cloning of a  
CD28 cDNA by a high efficiency COS cell expression  
system. PNAS, 84, 8573-8577.
- 15 Balasubramaniam, A., Sheriff, S., Johnson, M.E.,  
Prabhakaran, M., Huang, Y., Fischer, J.E., and Chance,  
W.T. (1994). [D-Trp<sup>32</sup>]Neuropeptide Y: A competitive  
antagonist of NPY in rat hypothalamus. J. Med. Chem. 37:  
20 311-315.
- Benson, G. Anthony, and Spillane, William J. (1980).  
Sulfamic Acid and Its N-Substituted Derivatives.  
25 Chem.Rev. 80: 151-186.
- Berrettini, W.H., Kaye, W.H., Gwirtsman, H., and  
Allbright, A. (1988). Cerebrospinal fluid peptide YY  
immunoreactivity in eating disorders. Neuropsychobiol 19:  
30 121-124.
- Bradford, M.M. (1976). A rapid and sensitive method for  
the quantitation of microgram quantities of protein  
utilizing the principle of protein-dye binding. Anal.  
35 Biochem. 72: 248-254.

Brady, L.S., Smith, M.A., Gold, P.W., and Herkenham, M. (1990). Altered expression of hypothalamic neuropeptide Y mRNAs in food-restricted and food-deprived rats. Neuroendocrinology 52: 441-447.

5

Chakraborty, T.K., Reddy, G.V., and Hussain, K. Azhar. (1991). Diastereoselective Strecker Synthesis using  $\alpha$ -Phenylglycinol as Chiral Auxiliary. Tetrahedron Letters. 32: 7597-7600.

10

Chance, W.T., Sheriff, S., Foley-Nelson, T., Fischer, J.E., and Balasubramaniam, A. (1989). Pertuss toxin inhibits neuropeptide Y-induced feeding in rats. Peptides 10, 1283-1286.

15

Clark, J.T. (1994). Aging-induced decrements in neuropeptide Y: The retention of ejaculatory behavior is associated with site-selective differences. Neurobiology of Aging 15: 191-196.

20

Clark, J.T., Kalra, P.S., Crowley, W.R., and Kalra, S.P. (1984). Neuropeptide Y and human pancreatic polypeptide stimulate feeding behavior in rats. Endocrinology 115: 427-429.

25

Cox, H., and Cuthbert, A.W. (1990). The effects of neuropeptide Y and its fragments upon basal and electrically stimulated ion secretion in rat jejunum mucosa. Br. J. Pharmac. 101: 247-252.

30

Cullen, B. (1987). Use of eukaryotic expression technology in the functional analysis of cloned genes. Methods Enzymol. 152: 685-704.

-121-

Dryden, S., Frankish, H., Wang, Q., and Williams, G. (1994). Neuropeptide Y and energy balance: one way ahead for the treatment of obesity? Eur. J. Clin. Invest. 24: 293-308.

5

Dumont, Y., J.-C. Martel, A. Fournier, S. St-Pierre, and R. Quirion. (1992). Neuropeptide Y and neuropeptide Y receptor subtypes in brain and peripheral tissues. Progress in Neurobiology 38: 125-167.

10

Dumont, Y., Fournier, A., St-Pierre, S., Quiron, R. (1995). Characterization of Neuropeptide Y Binding Sites in Rat Brain Membrane Preparations Using [<sup>125</sup>I][Leu<sup>31</sup>,Pro<sup>34</sup>] Peptide YY and [<sup>125</sup>I]Peptide YY<sub>1-36</sub> as Selective Y<sub>1</sub> and Y<sub>2</sub> Radioligands. J. Pharm. Exper. Ther. 272:(2) 673-680.

15

Eva, C., Oberto, A., Sprengel, R. and E. Genazzani. (1992). The murine NPY-1 receptor gene: structure and delineation of tissue specific expression. FEBS lett. 314: 285-288.

20

Eva, C., Keinänen, K., Monyer, H., Seeburg, P., and Sprengel, R. (1990). Molecular cloning of a novel G protein-coupled receptor that may belong to the neuropeptide receptor family. FEBS Lett. 271, 80-84.

25

Fuhlendorff, J., U. Gether, L. Aakerlund, N. Langeland-Johansen, H. Thøgersen, S.G. Melberg, U. B. Olsen, O. Thastrup, and T.W. Schwartz. (1990). [Leu<sup>31</sup>,Pro<sup>34</sup>]Neuropeptide Y: A specific Y<sub>1</sub> receptor agonist. Proc. Natl. Acad. Sci. USA 87: 182-186.

30

Gerald, C., Adham, A., Kao, HT, Olsen, M.A., Laz, T.M., Vaysse, P., Hartig, P.R., Branchek, T.A. and R.L. Weinshank. The 5-HT<sub>4</sub> receptor: molecular cloning and

35

-122-

pharmacological characterization of two splice variants  
(submitted for publication).

- 5 Grouzman, E., Comoy, E., and Bohuon, C. (1989). Plasma  
neuropeptide Y concentrations in patients with  
neuroendocrine tumors. J. Clin. Endoc. Metab. 68: 808-  
813.
- 10 Grundemar, L. and Rl Hakanson (1994). Neuropeptide Y  
effector systems: perspectives for drug development.  
Trends. Pharmacol. 15:153-159.
- 15 Grundemar, L., J.L. Krstenansky, and R. Hakanson. (1992).  
Activation of neuropeptide Y1 and neuropeptide Y2  
receptors by substituted and truncated neuropeptide Y  
analogs: identification of signal epitopes. Eur. J.  
Pharmacol. 232: 271-278.
- 20 Gubler, U abd B.J. Hoffman. (1983). A simple and very  
efficient method for generating cDNA libraries. Gene.  
25, 263-269
- 25 Hau, X.-Y., Boublik, J.H., Spicer, M.A., Rivier, J.E.,  
Brown, M.R., and Yaksh, T.L. (1991). The antinociceptive  
effects of spinally administered neuropeptide Y in the  
rat: Systematic studies on structure-activity  
relationship. JPET 258: 243-253.
- 30 Hauser, G.J., Myers, A.K., Dayao, E.K., and Zukowska-  
Grojec, Z. (1993). Neuropeptide Y infusion improves  
hemodynamics and survival in rat endotoxic shock. Am. J.  
Physiol. 265: H1416-H1423.
- 35 Heilig, M., and Widerlov, E. (1990). Neuropeptide Y: an  
overview of central distribution, functional aspects, and

-123-

possible involvement in neuropsychiatric illnesses. Acta Psychiatr. Scand. 82: 95-114.

5 Herzog, H., Y.J. Hort, H.J. Ball, G. Hayes, J. Shine, and  
L. Selbie. (1992). Cloned human neuropeptide Y receptor  
couples to two different second messenger systems. Proc.  
Natl. Acad. Sci. USA 89: 5794-5798.

10 Herzog, H., Y.J. Hort, H.J. Ball, G. Hayes, J. Shine, and  
L. Selbie. (1992). Cloned human neuropeptide Y receptor  
couples to two different second messenger systems. Proc.  
Natl. Acad. Sci. USA 89, 5794-5798.

15 Horstman, D.A., Brandon, S., Wilson A.L., Guyer, C.A.,  
Cragoe, E.J., Jr., Limbird, L.E. (1990). An Aspartate  
Conserved Among G-protein Receptors Confers Allosteric  
Regulation of Alpha2-Adrenergic Receptors by Sodium. J.  
Biol. Chem. 265: (35) 21590-21595.

20 Kalra, S.P., Fuentes, M., Fournier, A., Parker, S.L., and  
Crowley, W.R. (1992). Involvement of the Y-1 receptor  
subtype in the regulation of luteinizing hormone  
secretion by neuropeptide Y in rats. Endocrinology 130:  
3323-3330.

25 Kalra, S.P., Dube, M.G., Fournier, A., and Kalra, P.S.  
(1991). Structure-function analysis of stimulation of  
food intake by neuropeptide Y: Effects of receptor  
agonists. Physiology & Behavior 50: 5-9.

30 Kaye, W.H., Berrettini, W., Gwirtsman, H., and George,  
D.T. (1990). Altered cerebrospinal fluid neuropeptide Y  
and peptide YY immunoreactivity in anorexia and bulimia  
nervosa. Arch. Gen. Psychiat. 47: 548-556.

35

- 5 Kieffer, B., Befort, K., Gaveriaux-Ruff, C. and Hirth, C.G. (1992). The  $\delta$ -opioid receptor: Isolation of a cDNA by expression cloning and pharmacological characterization. Proc. natl. Acad. Sci. USA 89, 12048-12052.
- 10 Kingston, R. E. (1987) in Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. & Struhl, K. (Eds), *Current Protocols in Molecular Biology*, John Wiley and Sons, N.Y., Vol. 1, pp. 4.2.3-4.2.4.
- 15 Kloeck, J.A. and Leschinsky, K.L. (1976). An Improved Synthesis of Sulfamoyl Chlorides. J. Org. Chem. 41.
- Kluxen, F.W., Bruns, C. and Lubbert H. (1992). Expression cloning of a rat brain somatostatin receptor cDNA. Proc. Natl. Acad. Sci. USA 89, 4618-4622.
- 20 Kornfeld, R. and Kornfeld, S. (1985). Assembly of asparagine linked oligosaccharides. Annu. Rev. Biochem. 54, 631-664.
- 25 Kozak, M. (1989). The scanning model for translation: an update. J. Cell Biol. 108, 229-241.
- Kozak, M. (1991). Structural features in eukaryotic mRNAs that modulate the initiation of translation. J. Biol. Chem. 266, 19867-19870.
- 30 Krapcho, A. Paul and Kuell, Christopher S. (1990). Mono-protected Diamines, N-tert-Butoxycarbonyl- $\alpha,\omega$ -Alkanediamines from  $\alpha,\omega$ -Alkanediamines. Synthetic Communications. 20: 2559-2564.
- 35

- Krause, J., C. Eva, P.H. Seeburg, and R. Sprengel. (1991). Neuropeptide Y<sub>1</sub> subtype pharmacology of a recombinantly expressed neuropeptide receptor. Mol. Pharmacol. 41: 817-821.
- 5 Lacroix, J.S., Stjarne, P., Angard, A., and Lundberg, M. (1988). Sympathetic vascular control of the pig nasal mucosa: reserpine-resistant, non-adrenergic nervous responses in relation to neuropeptide Y and ATP. Acta  
10 Physiol. Scand. 133: 183-197.
- Landschultz, W.H., Johnson, P.F. and S.L. McKnight. (1988). The leucine zipper: a hypothetical structure common to a new class of DNA binding proteins. Science  
15 240, 1759-1764.
- Larhammar, D., A.G. Blomqvist, F. Yee, E. Jazin, H. Yoo, and C. Wahlestedt. (1992). Cloning and functional expression of a human neuropeptide Y/peptide YY receptor of the Y<sub>1</sub> type. J. Biol. Chem. 267: 10935-10938.  
20
- Levine, A.S., and Morley, J.E. (1984). Neuropeptide Y: A potent inducer of consummatory behavior in rats. Peptides 5: 1025-1029.  
25
- Maggio, R., Vogel Z. and J. Wess. (1993). Coexpression studies with mutant muscarinic/adrenergic receptors provide evidence for intermolecular "cross-talk" between G-protein-linked receptors. Proc. Natl. Acad. Sci. USA  
30 90: 3103-3107.
- Malaisse-Lagai, F., Carpentier, J.-L., Patel, Y.C., Malaisse, W.J., and Orci, L. (1977). Pancreatic polypeptide: A possible role in the regulation of food  
35 intake in the mouse. Hypothesis. Experientia 33: 915-917.

-126-

- McCormick, M. (1987). Sib Selection. Methods in Enzymology, 151: 445-449.
- 5 Miller, J. and Germain, R.N. (1986). Efficient cell surface expression of class II MHC molecules in the absence of associated invariant chain. J. Exp. Med. 164: 1478-1489.
- 10 Michel, M.C. (1991). Receptors for neuropeptide Y: multiple subtypes and multiple second messengers. Trends Pharmacol.: 12: 389-394.
- 15 Morley, J.E., and Flood, J.F. (1991). Neuropeptide Y and memory processing. An. N.Y. Acad. Sci. 611: 226-231.
- Okayama, H. and P. Berg (1983). A cDNA cloning vector that permits expression of cDNA inserts in mammalian cells. Mol. Cell. Biol. 3: 280-289.
- 20 O' Shea, R.D., and Gundlach, A.L. (1991). Preproneuropeptide Y messenger ribonucleic acid in the hypothalamic arcuate nucleus of the rat is increased in food deprivation or dehydration. J. Neuroendocrinol. 3: 11-14.
- 25 O'Shea, E.K., Rutkowski, R. and P.S. Kim. (1989). Evidence that the leucine zipper is a coiled coil. Science 243: 538-542.
- 30 Penner, S.B., Smyth, D.D., and Glavin, G.B. (1993). Effects of neuropeptide Y and [Leu<sup>31</sup>,Pro<sup>34</sup>]Neuropeptide Y on experimental gastric lesion formation and gastric secretion in the rat. JPET. 266: 339-343.



-127-

- Probst, W.C., Snyder, L.A., Schuster, D.I., Brosius, J and Sealfon, S.C. (1992). Sequence alignment of the G-protein coupled receptor superfamily. DNA and Cell Bio. 11, 1-20.
- 5 Sahu, A., and Kalra, S.P. (1993). Neuropeptidergic regulation of feeding behavior (neuropeptide Y). Trends Endocrinol. Metab. 4: 217-224.
- 10 Sato, M., Kawashima, Y., Goto, J., Yamane, Y., Chiba, Y., Jinno, S., Satake, M., Iwata, C. (1995). Synthesis and evaluation of novel fluorinated sulotroban-related sulfonamide derivatives as thromboxane A<sub>2</sub> receptor antagonists. Eur.J Med Chem. 30: 403-414.
- 15 Still, W. Clark, Kahn, Michael and Mitra, Abhijit. (1978). Rapid Chromatographic Technique for Preparative Separations with Moderate Resolution. J.Org.Chem. 43: 2923.
- 20 Robert, J.J., Grosco, M., Rouch, C., Jacquot, C., Cohen, Y. (1989) Unexpected Responses of the Obese "Cafeteria" Rat to the Peptide FMRF-Amide. Pharm. Bioch. Behavior 34: 341-344.
- 25 Rizzi, M., Samini, R., Sperk, G., and Vezzani, A. (1993). Electrical kindling of the hippocampus is associated with functional activation of neuropeptide Y-containing neurons. Eur. J. Neuroscience 5: 1534-1538.
- 30 Sanacora, G., Kershaw, M., Finkelstein, J.A., and White, J.D. Increased hypothalamic content of preproneuropeptide Y messenger ribonucleic acid in genetically obese Zucker rats and its regulation by food deprivation.
- 35 Endocrinology 127: 730-737 (1990).

Schwartz, T.W., J. Fuhlendorff, L.L.Kjems, M.S. Kristensen, M. Vervelde, M. O'Hare, J.L. Krstenansky, and B. Bjornholm. (1990). Signal epitopes in the three-dimensional structure of neuropeptide Y. Ann. N.Y. Acad. Sci. 611: 35-47.

Stanley, B.G., Magdalin, W., Seirafi, A., Nguyen, M.M., and Leibowitz, S.F. (1992). Evidence for neuropeptide Y mediation of eating produced by food deprivation and for a variant of the Y<sub>1</sub> receptor mediating this peptide's effect. Peptides 13: 581-587.

Stanley, B.G., and Leibowitz, S.F. (1984). Neuropeptide Y: Stimulation of feeding and drinking by injection into the paraventricular nucleus. Life Sci. 35: 2635-2642.

Stanley, B. G. Neuropeptide Y in multiple hypothalamic sites controls eating behavior, endocrine, and autonomic systems for body energy balance. In: *The Biology of Neuropeptide Y and Related Peptides*, pp. 457-509. Eds. W.F. Colmers and C. Wahlestedt. Humana Press, Totowa, New Jersey (1993).

Stanley, B.G., Daniel, D.R., Chin, A.S., and Leibowitz, S.F. (1985). Paraventricular nucleus injections of peptide YY and neuropeptide Y preferentially enhance carbohydrate ingestion. Peptides 6: 1205-1211.

Wahlestedt, C., L. Edvinsson, E. Ekblad, and R. Hakanson. Effects of neuropeptide Y at sympathetic neuroeffector junctions: Existence of Y<sub>1</sub> and Y<sub>2</sub> receptors. In: *Neuronal messengers in vascular function*, Fernstrom Symp. No 10., pp. 231-242. Eds A. Nobin and C.H. Owman. Elsevier: Amsterdam (1987).

-129-

Wahlestedt, C., Karoum, F., Jaskiw, G., Wyatt, R.J.,  
Larhammar, D., Ekman, R., and Reis, D.J. (1991). Cocaine-  
induced reduction of brain neuropeptide Y synthesis  
dependent on medial prefrontal cortex. Proc. Natl. Acad.  
5 Sci. 88: 2978-2082.

Wahlestedt, C., Regunathan, S., and D.J. Reis (1991).  
Identification of cultured cells selectively expressing  
Y1-, Y2-, or Y3-type receptors for neuropeptide Y/peptide  
10 YY. Life Sciences 50: PL-7 - PL-12.

Wahlestedt, C., Pich, E.M., Koob, G.F., Yee, F., and  
Heilig, M. (1993). Modulation of anxiety and neuropeptide  
Y-Y1 receptors by antisense oligodeoxynucleotides.  
15 Science 259: 528-531.

Wahlestedt, C., and D.J. Reis. (1993). Neuropeptide Y-  
Related Peptides and Their Receptors--Are the Receptors  
Potential Therapeutic Targets? Ann. Rev. Pharmacol. Tox.  
20 32: 309-352

Warden, D. and H.V. Thorne. (1968). Infectivity of  
polyoma virus DNA for mouse embryo cells in presence of  
diethylaminoethyl-dextran. J. Gen. Virol. 3, 371.  
25

Wells, J.A. (1994). Structural and functional basis for  
hormone binding and receptor oligomerization. Current  
Opinion in Cell Biology 6: 163-173

30 White, J.D., Olchovsky, D., Kershaw, M., and Berelowitz,  
M. (1990). Increased hypothalamic content of  
preproneuropeptide-Y messenger ribonucleic acid in  
streptozotocin-diabetic rats. Endocrinology 126: 765-772.

Zoubek, J., Somogyi, G.T., and De Groat, W.C. (1993). A comparison of inhibitory effects of neuropeptide Y on rat urinary bladder, urethra, and vas deferens. *Am. J. Physiol.* 265: R536-R543.

5

Zukowska-Grojec, Z., Haass, M., and Bayorh, M. (1986). Neuropeptide Y and peptide YY mediate non-adrenergic vasoconstriction and modulate sympathetic responses in rats. Reg. Pept. 15: 99-110.

10

Zukowska-Grojec, Z., Bergeson, S., Kuch-Wocial, A., and Colton, C. (1993). Mitogenic effect of neuropeptide Y in rat vascular smooth muscle cells. Neuropeptide Y Conference Abstracts, (Cambridge) C10.

15


-131-

5


What is claimed is:

1. A compound having the structure:

10



wherein Ar is



15

wherein each Z is independently N or C;

wherein each Y is independently N or C;

20

wherein p is an integer from 0 to 2;

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

-132-

wherein V is S, O, N, or  $\text{NR}_5$ ;

wherein X is a single bond or  $-\text{NH}-$ ;


- 5 wherein each  $\text{R}_2$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;
- 10 phenoxy; phenyl; pyridyl; thiophenyl; naphthyl; phthalimide;  $\text{C}_5\text{-C}_7$  lactam,  $\text{C}_5\text{-C}_7$  cyclic imide,  $\text{C}_5\text{-C}_7$  cyclic amino; wherein the phthalimide, lactam, cyclic imide, or cyclic amine is linked by nitrogen; and wherein the phenoxy, phenyl, pyridyl,
- 15 thiophenyl, naphthyl, phthalimide, lactam, cyclic imide, or cyclic amine is substituted with H, F, Cl, Br, I,  $\text{CF}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{C}_1\text{-C}_4$  alkylthio, or  $\text{NO}_2$ ;
- 20 wherein each  $\text{R}_3$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;
- 25 or  $\text{R}_2$  and  $\text{R}_3$  present on adjacent carbon atoms can constitute  $\text{C}_5\text{-C}_7$  cycloalkyl,  $\text{C}_5\text{-C}_7$  heterocycloalkyl or  $\text{C}_5\text{-C}_7$  heteroaryl;
- 30 wherein each  $\text{R}_4$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ; or  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;
- 35

-133-

wherein each  $R_5$  is independently H;  $C_1-C_3$  alkyl;  $C_2-C_6$  monohaloalkyl; or  $C_1-C_6$  polyhaloalkyl;

wherein  $L'$  is  $-NR_1-L-$  or


5




10

wherein  $L$  is  $C_3-C_6$  alkyl;  $C_3-C_9$  alkenyl;  $C_3-C_9$  alkynyl;


15



20



25




30

or

5

10



15

wherein  $R_1$  is H; or  $C_1$ - $C_6$  straight chained alkyl;

wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

20

wherein one dashed line is a double bond and the other dashed line is a single bond;

25

wherein each  $R_6$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_6$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_5$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

30

wherein i is an integer from 1 to 4; wherein n is an integer from 0 to 3; wherein m is an integer from 0 to 3;




-135-


wherein K is  $-\text{CH}_2-\text{NR}_{10}-\text{CHR}_7-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NR}_{10}-\text{CO}-(\text{CH}_2)_j-$   
 ;  $-\text{CH}_2-\text{NH}-\text{CO}-\text{NH}-(\text{CH}_2)_j-$ ;  $-\text{CO}-\text{NH}-\text{CHR}_7-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NR}_{10}-$   
 $\text{CO}-\text{CHR}_7-(\text{CH}_2)_j$ ;  $-\text{CH}_2-\text{NR}_{10}-\text{CS}-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NH}-\text{CS}-\text{NH}-$   
 $(\text{CH}_2)_1-$ ;  $-\text{CS}-\text{NH}-\text{CHR}_7-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NR}_{10}-\text{CS}-\text{CHR}_7-(\text{CH}_2)_j$ ;  
 5 or  $-\text{CH}_2-\text{N}=\text{CSR}_1-\text{NH}-(\text{CH}_2)_j$ ;

wherein j is an integer from 0 to 3;

wherein  $R_7$  is H;  $\text{C}_1-\text{C}_6$  alkyl;  $\text{CH}_2\text{OR}_5$ ;  $(\text{CH}_2)_p\text{NHCO}_2\text{R}_5$ ;  
 10  $(\text{CH}_2)_p\text{NHSO}_2\text{R}_5$ ;  $\text{CH}_2\text{N}(\text{R}_{11})_2$ ; phenyl; pyridyl; thiophenyl;  
 or naphthyl;

wherein W is





wherein Q is O; S; N;  $NR_4$ ; or  $C(R_5)_2$ ;

5

wherein b is an integer from 1 to 2;

10

wherein  $R_6$  is independently H; F; Cl; Br; I;  $NO_2$ ; OH;  $=O$ ;  $C_1-C_4$  alkyl;  $C_1-C_4$  alkoxy;  $C_1-C_4$  hydroxyalkyl;  $C_1-C_4$  methoxyalkyl;  $C_1-C_4$  monohaloalkyl;  $C_1-C_4$  polyhaloalkyl;  $N(R_5)_2$ ;  $NHCO_2R_4$ ;  $N(COR_5)_2$ ;  $NHCO_2R_4$ ;  $NHCONHR_5$ ;  $NHSO_2R_5$ ;  $N(SO_2R_5)_2$ ;  $CO_2R_5$ ;  $CON(R_5)_2$ ; or  $SO_2N(R_5)_2$ ;

15

wherein  $R_4$  is H;  $C_1-C_3$  alkyl;  $COR_5$ ;  $CO_2R_5$ ;  $CON(R_5)_2$ ;

wherein  $R_{10}$  is H; or  $C_1-C_6$  alkyl;

wherein  $R_{11}$  is H;  $COR_5$ ;  $COR_{12}$ ;  $SO_2R_5$ ;  $SO_2R_{12}$ ; and

wherein  $R_{12}$  is phenoxy; phenyl, pyridyl; thiophenyl;  
 or naphthyl; wherein the phenoxy, phenyl, pyridyl,  
 thiophenyl or naphthyl is substituted with H, F, Cl,  
 Br, I,  $CF_3$ ,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$  alkylthio,  
 $NO_2$ , phenyl, pyridyl or thiophenyl;

or a pharmaceutically acceptable salt thereof.

2. An (+) enantiomer of the compound of claim 1.

3. An (-) enantiomer of the compound of claim 1.






4. A compound of claim 1, wherein  $R_1$  is H;

wherein L is selected from  $C_3-C_9$  alkyl or




wherein the alkyl is substituted with H,  $OR_5$ , CN,  
 $C_1-C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl,  
 thiophenyl or naphthyl; and

wherein W is



5. A compound of claim 4, wherein Ar is selected from:



15

wherein each of  $R_2$ ,  $R_3$  and  $R_4$  is independently H; F, Cl, Br or I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl; or  $\text{N}(\text{R}_5)_2$ ;


20

wherein X is a single bond;

wherein each  $R_5$  is independently  $\text{C}_1\text{-C}_4$  alkyl;

wherein L is selected from  $\text{C}_3\text{-alkyl}$  or  $\text{C}_7\text{-alkyl}$ ;

25




or

30

35


wherein  $R_7$  is H;  $\text{CH}_2\text{OH}$ ; or  $\text{CH}_2\text{OR}_8$ ;

wherein W is



and wherein  $R_9$  is H; or  $C_1$ - $C_4$  alkyl.


- 5      6. A compound of claim 5, wherein Ar is selected from:



10

7. A compound of claim 6, wherein L is




15



-142-

8. A compound of claim 7, wherein K is  
-CH<sub>2</sub>-NR<sub>10</sub>-CHR<sub>7</sub>-(CH<sub>2</sub>)<sub>3</sub>-.


9. The compound of claim 8, wherein the compound is  
selected from the group consisting of:




35




5



10




15



20


25




30

35

5




10




15

20




25

30



35


5



10

or

15




20

10. A compound of claim 6, wherein L is C<sub>5</sub>-alkyl or C<sub>7</sub>-alkyl.

11. A compound of claim 10 having the structure:


25



30


or

35




12. A compound of claim 10, wherein K is  
 5 -CH<sub>2</sub>-NR<sub>10</sub>-CO-(CH<sub>2</sub>)<sub>3</sub>-.

13. A compound of claim 12 having the structure:



14. A compound of claim 10, wherein K is  
 15 -CH<sub>2</sub>-NH-CO-NH-(CH<sub>2</sub>)<sub>3</sub>-.

15. The compound of claim 14 having the structure:




16. A method of modifying feeding behavior of a subject  
 30 which comprises administering to the subject an

-147-


amount of a compound effective to decrease the consumption of food by the subject so as to thereby modify feeding behavior of the subject, wherein the compound has the structure:

5



10

wherein Ar is



5

wherein each Z is independently N or C;

wherein each Y is independently N or C;

10

wherein p is an integer from 0 to 2;

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

15

wherein V is S, O, N, or  $NR_5$ ;

wherein X is a single bond or  $-NH-$ ;

5 wherein each  $R_2$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; phenoxy; phenyl; pyridyl; thiophenyl; naphthyl; phthalimide;  $\text{C}_5\text{-C}_7$  lactam,  $\text{C}_5\text{-C}_7$  cyclic imide,  $\text{C}_1\text{-C}_7$  cyclic amino; wherein the phthalimide, lactam, cyclic imide, or cyclic amine is linked by nitrogen; and wherein the phenoxy, phenyl, pyridyl, thiophenyl, naphthyl, phthalimide, lactam, cyclic imide, or cyclic amine is substituted with H, F, Cl, Br, I,  $\text{CF}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{C}_1\text{-C}_4$  alkylthio, or  $\text{NO}_2$ ;


15 wherein each  $R_3$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_2\text{-C}_4$  alkenyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ;  $\text{SO}_2\text{N}(\text{R}_5)_2$ ; or  $R_2$  and  $R_1$  present on adjacent carbon atoms can constitute  $\text{C}_5\text{-C}_7$  cycloalkyl,  $\text{C}_5\text{-C}_7$  heterocycloalkyl or  $\text{C}_5\text{-C}_7$  heteroaryl;

25 wherein each  $R_4$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NHSO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ; or  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;




30 wherein each  $R_5$  is independently H;  $\text{C}_1\text{-C}_3$  alkyl;  $\text{C}_1\text{-C}_3$  monohaloalkyl; or  $\text{C}_1\text{-C}_3$  polyhaloalkyl;

35 wherein  $L'$  is  $-\text{NR}_1\text{-L-}$  or

-150-




wherein L is C<sub>3</sub>-C<sub>9</sub> alkyl; C<sub>3</sub>-C<sub>9</sub> alkenyl; C<sub>3</sub>-C<sub>9</sub> alkynyl;



or



-151-



wherein  $R_1$  is H; or  $C_1$ - $C_3$  straight chained alkyl;

15 wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

wherein one dashed line is a double bond and the other dashed line is a single bond;

20 wherein each  $R_i$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_6$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_5$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

25

wherein  $i$  is an integer from 1 to 4; wherein  $n$  is an integer from 0 to 3; wherein  $m$  is an integer from 0 to 3;

30

wherein  $K$  is  $-CH_2-NR_{10}-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-(CH_2)_3-$ ;  $-CH_2-NH-CO-NH-(CH_2)_3-$ ;  $-CO-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-(CH_2)_3-$ ;  $-CH_2-NH-CS-NH-$

-152-


$(\text{CH}_2)_1-$ ;  $-\text{CS}-\text{NH}-\text{CHR}_7-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NR}_{10}-\text{CS}-\text{CHR}_7-(\text{CH}_2)_j$ ;  
or  $-\text{CH}_2-\text{N}=\text{CSR}_1-\text{NH}-(\text{CH}_2)_j$

wherein  $j$  is an integer from 0 to 3;

5

wherein  $R_7$  is H;  $\text{C}_1$ - $\text{C}_6$  alkyl;  $\text{CH}_2\text{OR}_5$ ;  $(\text{CH}_2)_p\text{NHCO}_2\text{R}_5$ ;  
 $(\text{CH}_2)_p\text{NHSO}_2\text{R}_5$ ;  $\text{CH}_2\text{N}(\text{R}_{11})_2$ ; phenyl; pyridyl; thiophenyl;  
or naphthyl;





wherein Q is O; S; N; NR<sub>4</sub>; or C(R<sub>5</sub>)<sub>2</sub>;

5

wherein b is an integer from 1 to 2;

10

wherein R<sub>6</sub> is independently H; F; Cl; Br; I; NO<sub>2</sub>; OH; =O; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>; NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;

15

wherein R<sub>9</sub> is H; C<sub>1</sub>-C<sub>4</sub> alkyl; COR<sub>5</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>;

wherein R<sub>10</sub> is H; or C<sub>1</sub>-C<sub>4</sub> alkyl;

wherein R<sub>11</sub> is H; COR<sub>5</sub>; COR<sub>12</sub>; SO<sub>2</sub>R<sub>5</sub>; SO<sub>2</sub>R<sub>12</sub>; and

-155-



wherein  $R_{12}$  is phenoxy; phenyl, pyridyl; thiophenyl;  
or naphthyl; wherein the phenoxy, phenyl, pyridyl,  
thiophenyl or naphthyl is substituted with H, F, Cl,  
Br, I,  $CF_3$ ,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$  alkylthio,  
5  $NO_2$ , phenyl, pyridyl or thiophenyl;

or a pharmaceutically acceptable salt thereof.

10 17. The method of claim 16, wherein the compound is  
administered in combination with food.


18. The method of claim 16, wherein the subject is a  
vertebrate, a mammal, a human or a canine.

15 19. The method of claim 16, wherein the compound has the  
structure:




35


5




10



15



20




25

30


35

- 157 -


5



10




15



20


25




30

35

5




10



15


20



20. A method of treating a feeding disorder in a subject which comprises administering to the subject an amount of a compound effective to decrease the consumption of food by the subject, wherein the compound has the structure:

25

30




wherein Ar is

35



-159-



wherein each Z is independently N or C;

wherein each Y is independently N or C;

5 wherein p is an integer from 0 to 2;

wherein o is an integer from 0 to 1 and a is an integer from 0 to 3;

10 wherein V is S, O, N, or  $NR_5$ ;

wherein X is a single bond or  $-NH-$ ;

15 wherein each  $R_2$  is independently H; F; Cl; Br; I;  $NO_2$ ; OH;  $C_1-C_4$  alkyl;  $C_2-C_4$  alkenyl;  $C_1-C_4$  alkoxy;  $C_1-C_4$  hydroxyalkyl;  $C_1-C_4$  methoxyalkyl;  $C_1-C_4$  monohaloalkyl;  $C_1-C_4$  polyhaloalkyl;  $N(R_5)_2$ ;  $NHCO_2R_5$ ;  $N(COR_5)_2$ ;  $NHCO_2R_5$ ;

-160-

NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;  
 phenoxy; phenyl; pyridyl; thiophenyl; naphthyl;  
 phthalimide; C<sub>5</sub>-C<sub>7</sub> lactam, C<sub>5</sub>-C<sub>7</sub> cyclic imide, C<sub>5</sub>-C<sub>7</sub>  
 cyclic amino; wherein the phthalimide, lactam,  
 5 cyclic imide, or cyclic amine is linked by nitrogen;  
 and wherein the phenoxy, phenyl, pyridyl,  
 thiophenyl, naphthyl, phthalimide, lactam, cyclic  
 imide, or cyclic amine is substituted with H, F, Cl,  
 Br, I, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio,  
 10 or NO<sub>2</sub>;


wherein each R<sub>3</sub> is independently H; F; Cl; Br; I;  
 NO<sub>2</sub>; OH; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>2</sub>-C<sub>4</sub> alkenyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub>  
 hydroxyalkyl; C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl;  
 15 C<sub>1</sub>-C<sub>4</sub> polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>;  
 NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;  
 or R<sub>2</sub> and R<sub>3</sub> present on adjacent carbon atoms can  
 constitute C<sub>5</sub>-C<sub>7</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> heterocycloalkyl or  
 C<sub>5</sub>-C<sub>7</sub> heteroaryl;

20 wherein each R<sub>4</sub> is independently H; F; Cl; Br; I;  
 NO<sub>2</sub>; OH; C<sub>1</sub>-C<sub>4</sub> alkyl; C<sub>1</sub>-C<sub>4</sub> alkoxy; C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl;  
 C<sub>1</sub>-C<sub>4</sub> methoxyalkyl; C<sub>1</sub>-C<sub>4</sub> monohaloalkyl; C<sub>1</sub>-C<sub>4</sub>  
 polyhaloalkyl; N(R<sub>5</sub>)<sub>2</sub>; NHCOR<sub>5</sub>; N(COR<sub>5</sub>)<sub>2</sub>; NHCO<sub>2</sub>R<sub>5</sub>;  
 25 NHCONHR<sub>5</sub>; NHSO<sub>2</sub>R<sub>5</sub>; N(SO<sub>2</sub>R<sub>5</sub>)<sub>2</sub>; CO<sub>2</sub>R<sub>5</sub>; CON(R<sub>5</sub>)<sub>2</sub>; or  
 SO<sub>2</sub>N(R<sub>5</sub>)<sub>2</sub>;

30 wherein each R<sub>6</sub> is independently H; C<sub>1</sub>-C<sub>3</sub> alkyl; C<sub>1</sub>-C<sub>3</sub>  
 monohaloalkyl; or C<sub>1</sub>-C<sub>3</sub> polyhaloalkyl;


35 wherein L' is -NR<sub>1</sub>-L- or

-161-




5


wherein L is C<sub>3</sub>-C<sub>9</sub> alkyl; C<sub>3</sub>-C<sub>9</sub> alkenyl; C<sub>3</sub>-C<sub>9</sub> alkynyl;



10



15



20


25

or

30

35

-162-



wherein  $R_1$  is H; or  $C_1$ - $C_3$  straight chained alkyl;

15 wherein the alkyl, alkenyl or alkynyl is substituted with H,  $OR_5$ , CN,  $C_1$ - $C_6$  alkyl,  $CH_2OR_5$ ,  $CON(R_5)_2$ ,  $CO_2R_5$ , phenyl, pyridyl, thiophenyl or naphthyl;

wherein one dashed line is a double bond and the other dashed line is a single bond;

20 wherein each  $R_6$  is independently H; CN;  $OR_5$ ;  $C_1$ - $C_6$  alkyl;  $CH_2OR_5$ ;  $CON(R_5)_2$ ;  $CO_2R_5$ ; phenyl; pyridyl; thiophenyl or naphthyl; wherein the phenyl, pyridyl, thiophenyl or naphthyl is substituted with H, F, Cl, Br, I,  $CF_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_1$ - $C_4$  alkylthio, or  $NO_2$ ;

25 wherein  $i$  is an integer from 1 to 4; wherein  $n$  is an integer from 0 to 3; wherein  $m$  is an integer from 0 to 3;

30 wherein  $K$  is  $-CH_2-NR_{10}-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-(CH_2)_3-$ ;  $-CH_2-NH-CO-NH-(CH_2)_3-$ ;  $-CO-NH-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CO-CHR_7-(CH_2)_3-$ ;  $-CH_2-NR_{10}-CS-(CH_2)_3-$ ;  $-CH_2-NH-CS-NH-$

-163-


$(\text{CH}_2)_1-$ ;  $-\text{CS}-\text{NH}-\text{CHR}_7-(\text{CH}_2)_j-$ ;  $-\text{CH}_2-\text{NR}_{10}-\text{CS}-\text{CHR}_7-(\text{CH}_2)_j$ ;  
or  $-\text{CH}_2-\text{N}=\text{CSR}_1-\text{NH}-(\text{CH}_2)_j$ ;


wherein  $j$  is an integer from 0 to 3;

5

wherein  $R_7$  is H;  $\text{C}_1-\text{C}_6$  alkyl;  $\text{CH}_2\text{OR}_5$ ;  $(\text{CH}_2)_p\text{NHCO}_2\text{R}_5$ ;  
 $(\text{CH}_2)_p\text{NHSO}_2\text{R}_5$ ;  $\text{CH}_2\text{N}(\text{R}_{11})_2$ ; phenyl; pyridyl; thiophenyl;  
or naphthyl;

wherein W is





wherein Q is O; S; N; NR<sub>9</sub>; or C(R<sub>5</sub>)<sub>2</sub>;

5            wherein b is an integer from 1 to 2;

10 wherein  $R_8$  is independently H; F; Cl; Br; I;  $\text{NO}_2$ ; OH;  $=\text{O}$ ;  $\text{C}_1\text{-C}_4$  alkyl;  $\text{C}_1\text{-C}_4$  alkoxy;  $\text{C}_1\text{-C}_4$  hydroxyalkyl;  $\text{C}_1\text{-C}_4$  methoxyalkyl;  $\text{C}_1\text{-C}_4$  monohaloalkyl;  $\text{C}_1\text{-C}_4$  polyhaloalkyl;  $\text{N}(\text{R}_5)_2$ ;  $\text{NHCOR}_5$ ;  $\text{N}(\text{COR}_5)_2$ ;  $\text{NHCO}_2\text{R}_5$ ;  $\text{NHCONHR}_5$ ;  $\text{NH}\text{SO}_2\text{R}_5$ ;  $\text{N}(\text{SO}_2\text{R}_5)_2$ ;  $\text{CO}_2\text{R}_5$ ;  $\text{CON}(\text{R}_5)_2$ ; or  $\text{SO}_2\text{N}(\text{R}_5)_2$ ;

wherein  $R_9$  is H;  $C_1$ - $C_3$  alkyl;  $COR_5$ ;  $CO_2R_5$ ;  $CON(R_5)_2$ ;

wherein R<sub>10</sub> is H; or C<sub>1</sub>-C<sub>6</sub> alkyl;

wherein  $R_{11}$  is H;  $COR_5$ ;  $COR_{12}$ ;  $SO_2R_5$ ;  $SO_2R_{12}$ ; and

-166-

5 wherein R<sub>12</sub> is phenoxy; phenyl, pyridyl; thiophenyl;  
or naphthyl; wherein the phenoxy, phenyl, pyridyl,  
thiophenyl or naphthyl is substituted with H, F, Cl,  
Br, I, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio,  
NO<sub>2</sub>, phenyl, pyridyl or thiophenyl;

or a pharmaceutically acceptable salt thereof.





10 21. The method of claim 20, wherein the feeding disorder  
is obesity or bulimia.

22. The method of claim 21, wherein the subject is a  
vertebrate, a mammal, a human or a canine.


15




23. The method of claim 20, wherein the compound has the structure:




5




10



15




20



25


30




35

-169-


5



10




15



20

or

25



30

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/19085**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 31/18, 31/335, 31/47; C07C 311/10; C07D 215/12

US CL : 514/311, 452, 602; 546/165, 172; 549/366; 564/87, 90, 92, 94

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/311, 452, 602; 546/165, 172; 549/366; 564/87, 90, 92, 94

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS Online structure

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A         | US 5,352,705 A (DEANNA, deceased et al.) 04 October 1994, see entire document.     | 1-23                  |
| A         | US 5,455,258 A (MACPHERSON et al.) 03 October 1995, see entire document.           | 1-23                  |

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

|   |  |
|---|--|
| * Special categories of cited documents:  | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  |
| *A* document defining the general state of the art which is not considered to be of particular relevance  | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone   |
| *E* earlier document published on or after the international filing date  | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *G* document member of the same patent family  |
| *O* document referring to an oral disclosure, use, exhibition or other means  |  |
| *P* document published prior to the international filing date but later than the priority date claimed  |  |

|   |  |
|---|--|
| Date of the actual completion of the international search<br>31 MARCH 1997  | Date of mailing of the international search report<br>05 MAY 1997                    |
| Name and mailing address of the ISA/US<br>Commissioner of Patents and Trademarks<br>Box PCT<br>Washington, D.C. 20231<br>Facsimile No. (703) 305-3230 | Authorized officer <i>IW for</i><br>JACQUELINE HALEY<br>Telephone No. (703) 308-1235 |